

# VISION GUIDELINES<sup>1</sup>

## I. INTRODUCTION

### A. OUTLINE OF HIGHLIGHTED CONDITIONS

- 1) Far Acuity Deficiency
  - Methods of correction
    - Spectacles
    - Contact lenses
    - Orthokeratology
- 2) Refractive Surgical Procedures
  - LASIK, LASEK, PRK, & SMILE
  - Phakic Intraocular Lenses
  - Radial Keratotomy
- 3) Color Vision Deficiency
- 4) Other Visual Functions
  - Visual Field Deficiency
  - Binocular Fusion Deficiency
  - Contrast Sensitivity

A summary of the recommended evaluation criteria presented in this chapter begins on page [XI-60](#).

### B. PRE-EMPLOYMENT VISION SCREENING AND THE LAW

Despite the importance of vision to the safety of the officer and the public, pre-employment vision standards have been the subject of several legal challenges. Most commonly, agency vision standards have been assailed for: (1) insufficient job-relatedness; (2) failure to allow for reasonable accommodation; (3) inconsistency in standards across agencies; and (4) inconsistent enforcement of standards within an agency, particularly with respect to candidates versus incumbents.

- 1) **Insufficient Job Relatedness.** An agency's selection of a vision standard must be based on job-relatedness rather than unsubstantiated suppositions. The vision guidelines presented here are supported by detailed, quantitative summaries of the currently available literature. *However, it is incumbent upon each agency to review to ensure that the assumptions and findings upon which these guidelines are based upon are sufficiently applicable to the job duties and circumstances in its own jurisdiction.*

---

<sup>1</sup> **Authors:** Jeffery Hovis, O.D., Ph.D. (MED-TOX) with assistance from R. Leonard Goldberg, M.D.  
**Specialist Review Panel:** James Bailey, O.D.; Gregory Good, O.D.; James Sheedy, Ph.D., O.D.; Melissa Barnett, O.D.

- 2) **Failure to Allow for Reasonable Accommodation and Mitigating Measures.** Another frequently adjudicated agency vision policy is the unilateral prohibition against the use of vision correction devices or procedures (e.g., spectacles, contact lenses, refractive surgery). Findings in favor of the candidate are not uncommon when the agency appears to have based its policy on unfounded concerns rather than factual evidence. Included in this section is a detailed discussion of the advantages and risks associated with each method of vision correction and resulting recommendations for establishing far acuity standards.
- 3) **Agency-Specific Job Demands and Conditions.** While patrol officers across the state share many essential job functions, differences in job demands and environmental conditions exist across agencies. Thus, the risk posed by an officer with decreased visual function (or the hardship caused by accommodating such individuals) may also vary across agencies. Throughout this section, the influence of position and agency-specific factors are discussed to enable the creation of vision standards that are appropriate for each department.
- 4) **Inconsistent Enforcement of Agency Standards.** An agency's allegation that its vision standards are job-related is weakened if incumbent officers who no longer meet these standards are successfully performing the job. While at times judges have agreed with law enforcement agency assertions that experience can partially compensate for visual impairment (e.g., *Padilla v. City of Topeka*, 1985), other courts have ruled against law enforcement agencies who maintain stringent vision standards for candidates while failing to enforce these standards among its incumbent officers (e.g., *Brown County v. LIRC*, 1985).

Fortunately, the stability of most visual functions renders this issue largely moot. Except for near vision, the visual acuity of the vast majority of persons remains stable with age. As evidence, the results of uncorrected vision testing among incumbents of the Los Angeles City Fire Department (Goldberg & Bible, 1993) showed that, after an average of 11 years of service, over 96% of the 1,111 firefighters tested still possessed uncorrected vision that met the pre-placement guideline of 20/40. Even in the class of Captain II, approximately 90% of the 164 incumbents still maintained 20/40 vision after an average of 23 years of service. Good et al. (1998) reported similar results for the Columbus, Ohio police department. Although the average number of years of service was not stated, 94% of incumbent officers met the 20/40 uncorrected acuity that had been in place for 16 years; only 0.9% had an uncorrected acuity worse than 20/63.

## C. IMPLICATIONS FOR JOB PERFORMANCE

In 1984, POST conducted a vision-oriented job analysis for the position of patrol officer (Briggs, 1984). After interviewing and observing officers in the field, a panel of vision experts developed a list of 17 relevant visual skills. The importance of these skills for patrol officer performance was then rated by 158 incumbent officers (average patrol experience = 5 years) who had been shown slides depicting and illustrating each of the 17 visual skills. The officers were also asked to provide detailed accounts of actual critical incidents based on their personal experiences. The officers produced 1,291 incidents, which involved at least one of the 17 visual skills. The results from both activities are reported in Table XI-1.

As indicated in Table XI-1, no visual skill was rated less than "important." Officers rated dark adaptation as the most important visual skill, followed by peripheral vision. The ability to identify objects was involved in the highest percentage of critical incidents (24.9%), followed by visual pursuit (21.1%), motion detection (17.9%), dynamic far acuity (15.6%), dark adaptation (15.5%), and peripheral vision (11.2%). These results confirm the importance of virtually every visual capacity in the safe performance of patrol officer duties.

**Table XI-1.** Patrol Officer Importance Ratings of 17 Visual Skills (Briggs, 1984)

| Visual Skill                      | Importance Rating* (N=158) | % of the 1,291 Critical Incidents in Which Skill Was Involved |
|-----------------------------------|----------------------------|---|
| Dark Adaptation                   | 4.50                       | 15.5  |
| Peripheral Vision                 | 4.34                       | 11.2  |
| Identify Objects                  | 4.29                       | 24.9  |
| Motion Detection                  | 4.13                       | 17.9  |
| Fine Details/Various Light Levels | 4.03                       | 9.1   |
| Pursuit                           | 3.95                       | 21.1  |
| Dynamic Near Acuity               | 3.93                       | 2.5   |
| Accommodation                     | 3.87                       | 4.3   |
| Dynamic Far Acuity                | 3.81                       | 15.6  |
| Depth Perception                  | 3.68                       | 6.8   |
| Light Adaptation                  | 3.63                       | 3.3   |
| Glare Recovery                    | 3.61                       | 1.1   |
| Glare Tolerance                   | 3.59                       | 9.8   |
| Identify Large Forms              | 3.54                       | 1.1   |
| Static Far Acuity                 | 3.54                       | 3.8   |
| Color Identification              | 3.53                       | 5.8   |
| Color Discrimination              | 3.30                       | 1.2   |

\*Rating scale values: 5 = critically important, 4 = very important, 3 = important, 2 = of some importance, 1 = of little importance

Similar results were found for Ontario police officers, who were asked to identify visual skills essential for successful performance in three frequently occurring scenarios: impaired drivers, domestic disturbances, and breaking and entering (Shaw & Gledhill, 1995). The results are summarized in Table XI-2. Identifying objects was a frequently-identified visual skill, followed by seeing in poor visibility conditions (e.g., at night) and peripheral vision. In contrast to the POST study, the frequency of needing color vision was comparable to that of visual acuity. (This difference may be due, in part, to the limited number of scenarios addressed).

**Table XI-2.** Visual skills identified by Ontario police officers as essential for responding to impaired drivers, domestic disturbances, and breaking and entering (Shaw & Gledhill, 1995)

| Visual Skill  | Frequency that skill was mentioned (N=114) |
|---|--|
| Color Vision<br>Scan for evidence of personal offense such as blood and bruises, changes in complexion, noting details (vehicles, clothes and general description) for evidence and reporting descriptions                | 28.1%                                      |
| Visual Acuity<br>See where people go, get descriptions and details for reporting and evidence, not trip over things, read license plates, see with cruiser lights flashing  | 28.1%                                      |
| Visual Acuity (at night and in poor visibility)<br>See in poor light, night vision, artificial light, see in shadows. See in poor visibility, bad weather conditions, poorly lit environments, smoked windows on vehicles | 9.6%                                       |
| Peripheral Vision   | 9.6%                                       |
| Visual Acuity and Peripheral Vision<br>Must have good general observation skills - aware of surroundings, multiple inputs, description of person  | 9.6%                                       |
| Depth Perception  | 2.6%                                       |

## II. MEDICAL EXAMINATION AND EVALUATION GUIDELINES

### A. GENERAL SCREENING RECOMMENDATIONS

#### 1) History:

Candidates should be questioned regarding the use of spectacles or contact lenses, visual loss, night blindness, refractive surgery and a history of eye diseases ([Medical History Statement, Form #2-252](#)). The candidates' driver licenses should be reviewed to determine if they have any driving restrictions or conditions.

#### 2) Routine Testing:

##### a. FAR ACUITY

##### ***Far acuity testing procedures:***

*NOTE: It is very important to use standardized charts and methods when measuring visual acuity. Non-standardized testing results in erroneous measurements and increased measurement variability.*

1. Use only charts that meet, or exceed, the ANSI Z80.21 (2020) standards. The Bailey-Lovie chart and the ETDRS chart - both original and revised - meet this standard (Ferris, et al., 1982). Both charts have the additional advantage that the number of

letters on each line of these charts is equal up to 20/200, thereby preventing candidates from inferring that the largest letter on the chart is an “E.”

2. Use only charts designed for 13 feet (ft) or 20 ft test distances. Individuals with low to moderate uncorrected myopia (i.e., nearsightedness) may pass at closer viewing distances. The 13 ft charts adjust the letter sizes so that the angular size of the letters viewed from 13 ft is equal to the angular size of the letters viewed from 20 ft; therefore, no additional conversion should be necessary.
3. The chart should have relatively even luminance (i.e., brightness) across its surface: luminance should be 160 cd/m<sup>2</sup> with an acceptable range of 80-320 cd/m<sup>2</sup>. Self-illuminated charts are available from various ophthalmic supply companies. Printed charts are acceptable if there is sufficient light falling on the chart so that the light reflected from the white background falls within the acceptable range.<sup>2</sup> The illumination on the chart should range from 270 lx to 1000lx, with 530 lx being equivalent to 160 cd/m<sup>2</sup>.

If an illuminance meter is unavailable, direct two 60-watt incandescent lamps (or lights with a lumen output of at least 800 lumens) toward the chart. Two lamps are necessary to maintain a uniform illumination across the chart. The lamps should be separated by 30”-36” and placed 24” from the chart.

Do not place printed charts in hallways with overhead light fixtures without measuring the light falling on the chart. The variation in the intensity distribution of the light leaving the light fixtures and placement of the chart relative to the fixtures may result in an unacceptable amount of light reaching the chart and/or nonuniformities of the light reflected from various regions of the chart. The variance from the mean illumination should be within  $\pm 25\%$  (ISO 8596:2009).

4. The candidate's eyes should be evaluated to ensure that contact lenses are not worn during uncorrected vision testing.
5. Testing should be performed with the candidate at the chart's viewing distance (13 ft or 20 ft). If the candidate cannot discern the top row of letters at this distance, the acuity should be recorded as worse than the value corresponding to the top row.
6. Conduct monocular testing before binocular testing.
7. Measure uncorrected acuities before corrected acuities.
8. An occluder should cover one eye while testing the other eye. The candidate can hold the occluder. The occluder can simply be an index card.
9. Candidates should be informed that they cannot squint during the testing. The tester should observe the candidate to ensure compliance.
10. Candidates should read at least one line in which they can identify all five letters. It can be helpful to instruct candidates to start with the smallest line of letters that they can read easily. They should proceed to successively smaller acuity lines until they cannot correctly identify any letters on a line. They should be encouraged to guess when

---

<sup>2</sup> Meters to measure the illumination falling on the charts (such as General Light Meter Model # DLM1337 are available from Home Depot and other suppliers for ~\$100.

letter recognition becomes difficult. It is helpful to have a score sheet that corresponds to the chart for recording accuracy.

11. Candidates should be given credit for each letter properly identified. Record the acuity line in which at least 3 of the 5 letters are correctly identified. Include a +/- notation to precisely convey the number of letters properly identified. For example, if the candidate read the entire 20/30 line and one additional letter on the 20/25 line, the acuity would be 20/30+1. Identifying all of the 20/30 line and 3 of the 5 letters on the 20/25 line would result in an acuity of 20/25-2. Since the charts have 5 letters per acuity line, the +/- value will never exceed a value of 2. Letters that are correctly identified on a smaller line compensate for letters missed on a larger line. For example, if a candidate reads 4 of 5 letters on the 20/30 line and 2 of 5 on the 20/25 line, the score would be 20/30+1. The measured acuity should be compared to the agency standard. For example, if a standard has been set at 20/40, a measured acuity of 20/40-1 is not sufficient.

## **Use of Vision Screeners**

Vision screeners are portable, self-contained units with internally calibrated light levels. While popular in occupational medicine, several factors can limit their validity. First, the varying visual acuity charts available poses an impediment to reliability. In particular, there are two types of Landolt ring charts (where the letter C is rotated in different orientations). One has multiple rings on the same line; the candidate's task is to identify the orientation of the break in the ring. The other type has four (4) rings, three of which are continuous, and one has a break in the ring. The candidate must identify which ring is open AND the orientation of the opening, since the candidate could correctly identify which one of the four rings is open 25% of the time by guessing. Since standard scoring sheets only denote which ring has the gap, the score sheet must be modified to include gap orientation.

Hovis & Ramaswamy (2006) assessed police cadets who met the 20/20 acuity requirement with the Optec 2000 using the 4-ring target. Their results suggest that this chart may overestimate the acuity for some individuals;<sup>3</sup> 8% did not meet the acuity standard of 20/20 using the Bailey-Lovie chart, although no one had acuity worse than 20/25.

The perceived proximity of the acuity chart can cause some candidates to change the focus of their natural lenses as if they are looking at a near object, inducing artificial myopia. This may be the reason behind the reported low specificity of vision screeners for police candidates. In reviewing police candidate records, Hovis (2008) found that 35% of candidates whose acuities were worse than 20/40 when measured with the Optec 2000<sup>3</sup> obtained this level of acuity or better when measured with a wall visual acuity chart. Therefore, assessing visual acuity with visual screening units is not recommended. If candidates are assessed with a vision screener, those who fail to meet an acuity requirement should be retested with a wall chart.

## **COLOR VISION**

All candidates should be administered the 4<sup>th</sup> edition of the Hardy Rand Rittler (HRR) pseudoisochromatic plate color vision test.

The test must be administered under proper illumination conditions. All color vision tests are designed to be used with a standard source of illumination, one approximating standard

---

<sup>3</sup> They were not required to identify the location of the gap.

illumination "C" of the CIE (International Commission on Illumination). Ordinary daylight fluorescent, LEDs, compact fluorescent lamps (CFL) or incandescent lamps should NOT be used. The standard illuminant should be the only source of illumination. There should be no other light source in the room; any window shades should be drawn.

The True Daylight Illuminator is a standalone unit, consisting of an easel for the test and a light fixture containing a Verilux True Color Light fluorescent tube (F15T8VLX).<sup>4</sup> This unit is sufficient for illuminating color vision tests (Milburn & Mertens, 1993).

Many fluorescent lamps and some LEDs lamps are advertised as "full-spectrum" or "natural daylight." These may be suitable substitutes provided that they have both a color rendering index of at least 90 and a correlated color temperature between 5300 to 8500 K (Dain et al., 1993; Hovis & Neumann, 1995, Dain et al., 2020). These lamps can be found in the "professional lamps" or "high color rendering" section of an online catalog.

Tinted lenses alter the standard illumination required for all color vision tests, thereby invalidating the results. Therefore, colored contact lenses or tinted spectacle lenses (such as the X-Chrom, X-Chrome, Colormax, Enchroma, Chromagen, Vino) should not be permitted. The cornea of each eye should be examined for unreported contact lens wear, especially densely tinted lenses such as the X-Chrome lens. Due to the difficulty of judging the density of any contact lens tint, all tinted contact lenses should be removed before testing.

### Instructions for Administration and Interpretation of the HRR test

1. Before administering the test, the candidate, test, and illuminant should be properly positioned. The candidate should be seated 75 cm (about 30 in) from the test. The plates should be supported and then tilted until they are perpendicular to the candidate's line of sight. The illuminant should be situated so that the illumination is direct and even, and is incident approximately 45° to the plates. It is desirable to have a small paintbrush available for use as a pointer or for tracing symbols on the plates.
2. Open the book to the first demonstration plate and instruct the candidate: *"I am going to show several plates that may contain simple colored figures. Here are examples of a circle and X."* Show the candidate the next plate and state, *"Here are examples of a triangle and X."* Show the third plate and state *"Here is a circle. There can be one or two figures, or (as you turn to 4<sup>th</sup> plate) no figure. You need to point to the location of each figure that you see with the paintbrush. If you are not sure, trace the figure with the paintbrush."*
3. The order of presentation (and/or orientation of the page) should be randomized to ensure that candidates are not advantaged by prior experience with the test. The viewing time for each plate should be set at approximately 3 seconds.
4. A correct response identifies **both the shape(s) and location** of each figure on each page. Mark each figure on the score sheet as correct or incorrect. Each missed figure on a plate counts as an error (Cole et al., 2006).
  - a) Any error on plates 5 and 6 constitutes a failure on HRR blue-yellow plates. Failure on the red-green screening plates is more than one error on plates 7-10. Note: these validated failure criteria are different from the test publisher instructions (Cole et al., 2006; Almustanyir, 2014).

---

<sup>4</sup> Available from Good-Lite, Elgin, IL

- b) The severity of the **red-green defect** is graded as very mild, mild, medium and severe, depending on whether the candidate sees the figures on plates 11-20:

Very mild = errors on screening plates only; no errors on plates 11-20  
Mild = any errors on plates 11-15; no errors on plates 16-20  
Medium = any errors on plates 16-18; no errors on plates 19 and 20  
Severe = any errors on plates 19 and 20

The severity of the **blue-yellow defect** is graded as mild, medium and severe:

Mild = errors only on plates 5 and 6  
Medium = any errors on plates 21 and 22  
Severe = errors on plates 23 and 24

- c) If the candidate falls into the medium category by just one error on plates 16-18, color vision should be assessed with one of the other tests described under [Color Vision Deficiency](#).<sup>5</sup>
- d) Identifying the screening figures correctly but making errors on the diagnostic plates should be treated as a fail; the candidate should be assessed with one of the other tests described under [Color Vision Deficiency](#).

## **BINOCULAR VISION – EXTRAOCULAR MUSCLE DYSFUNCTION**

Candidates should be tested for gaze restrictions, noticeable strabismus and potential double vision as discussed in the physical examination of the eyes in [Chapter IX – Neurology](#). A history of strabismus surgery, eye patching, visual training, or prismatic corrections in spectacle lenses are also risk factors for double vision; these candidates should be evaluated by a vision specialist to determine the risk for double vision when tired or in a reduced visual environment.

## **B. EVALUATION OF COMMON CLINICAL SYNDROMES**

### **1) FAR ACUITY DEFICIENCY**

#### **a. GENERAL CONSIDERATIONS**

Various methods have been used to determine the impact of far acuity deficiencies on performance as a patrol officer. The critical patrol officer functions studied include: (1) deciding whether to discharge a firearm; (2) facial recognition; (3) license plate identification; and (4) withdrawal under fire.

#### **1. "Shoot-No-Shoot" Decisions**

Deciding whether to discharge a firearm is a critical, not-infrequent task facing patrol officers. In 1986, approximately 1/50 LAPD sworn officers discharged their weapon; 42% of these incidents resulted in a civilian being wounded or killed (Pate & Hamilton, 1991). Since this study included

---

<sup>5</sup>The additional test results should be used to determine whether the candidate has very mild-to-mild color vision deficiency or moderate-to-severe based on all available test results and history. Individuals with a moderate to severe color vision defect are disqualified.



officers who do not work in the field, the firearm discharge rate among officers assigned to field duty would be expected to be higher.

A separate study of LAPD officer-initiated shootings during 1990-92 found that over 30% of the 519 incidents occurring during this period involved shooting at targets over 25 ft away. Moreover, 65% of officer-initiated shootings took place at night or at dawn/dusk (Spilberg, 1993).

In 2010, the U.S. Department of Justice surveyed 295 officers who averaged 17 years of law enforcement experience. Of these officers, 96% responded that they drew their firearms at least once each year, and 59 (20%) had been involved in at least one critical incident where they had fired their weapon (Pinizzotto, 2012).

An officer's ability to rapidly determine whether a suspect is holding a weapon is typically studied using decorection lenses in scenarios at distances varying from 7-25 yards. In a 1981 study, six California Highway Patrol (CHP) officers with 20/20 or better uncorrected vision were sequentially decorected to 20/40, 20/80, and 20/200. During each visual condition, the officers were asked to identify whether a "suspect" was holding a gun or a comb at distances of 7, 15, and 25 yards. The study was conducted during the day under clear skies. No errors were made with 20/20 vision, even at a distance of 25 yards (Table XI-3). With 20/40 vision, the officers correctly identified all of the objects at 7 yards but misidentified 14% at 15 yards. With 20/80 vision, officers misidentified 8% of the objects at 7 yards and 22% of the objects at 15 yards (Giannoni, 1981).

**Table XI-3.** Percentage Correct Identifications for "Shoot" and "No Shoot" Scenario

| Candidates     | 25 Yard Distance |       |       |        | 15 Yard Distance |       |       |        | 7 Yard Distance |       |       |        | Combined Distances |       |       |        |
|----------------|------------------|-------|-------|--------|------------------|-------|-------|--------|-----------------|-------|-------|--------|--------------------|-------|-------|--------|
|                | 20/20            | 20/40 | 20/80 | 20/200 | 20/20            | 20/40 | 20/80 | 20/200 | 20/20           | 20/40 | 20/80 | 20/200 | 20/20              | 20/40 | 20/80 | 20/200 |
| <b>Cell B</b>  |                  |       |       |        |                  |       |       |        |                 |       |       |        |                    |       |       |        |
| 1              | 100              | 50.0  | 50.0  | 50.0   | 100              | 50.0  | 50.0  | 50.0   | 100             | 100   | 50.0  | 50.0   | 100                | 66.7  | 50.0  | 50.0   |
| 2              | 100              | 50.0  | 50.0  | 66.7   | 100              | 83.3  | 50.0  | 66.7   | 100             | 100   | 100.0 | 100.0  | 100                | 77.8  | 66.7  | 77.8   |
| 3              | 100              | 100.0 | 50.0  | 50.0   | 100              | 83.3  | 83.3  | 66.7   | 100             | 100   | 100.0 | 66.7   | 100                | 94.4  | 77.8  | 61.1   |
| <b>Cell A</b>  |                  |       |       |        |                  |       |       |        |                 |       |       |        |                    |       |       |        |
| 4              | 100              | 100.0 | 16.7  | 50.0   | 100              | 100.0 | 83.3  | 83.3   | 100             | 100   | 100.0 | 100.0  | 100                | 100.0 | 66.7  | 77.8   |
| 5              | 100              | 100.0 | 83.3  | 50.0   | 100              | 100.0 | 100.0 | 66.7   | 100             | 100   | 100.0 | 83.3   | 100                | 100.0 | 94.4  | 66.7   |
| 6              | 100              | 100.0 | 50.0  | 33.3   | 100              | 100.0 | 100.0 | 33.3   | 100             | 100   | 100.0 | 100.0  | 100                | 100.0 | 83.3  | 55.5   |
| <b>Average</b> | 100              | 83.3  | 50.0  | 50.0   | 100              | 86.1  | 77.8  | 61.1   | 100             | 100   | 91.7  | 83.3   | 100                | 89.8  | 73.2  | 64.8   |


From Giannoni, B. Entry-level vision requirements validation study. Personnel Bureau, California Highway Patrol. October 1981

This study was partially replicated in 1999 using a handgun and four different distracters: a cell phone, a brown beer bottle, a pair of aviator-style black sunglasses in the daytime trials and a purse at night (Carmean et al., 2000). Five university students participated in the experiment, which was conducted during the day (7,275 lx) and night (2 lx)<sup>6</sup>.

<sup>6</sup> Assuming that average reflectance of the objects in the scene was 0.30 and all objects were matte surfaces, the average luminance of the scene would be 695 cd/m<sup>2</sup> in the daytime and 0.2 cd/m<sup>2</sup> at night.

Figure XI-1 displays the percentage of participants who were able to identify all five objects correctly at each distance and level of visual acuity. All participants were able to identify all the objects correctly at 15 yards in the daytime at all levels of acuity; therefore, no further trials were conducted at the closer viewing distances for this condition. Similar to the Giannoni study, the results indicate that visual acuity must be better than 20/40 to distinguish a gun from other handheld items from a distance of 25 yards during daytime. The results also show how low light levels can affect performance. If the acuity (measured in the clinic) is 20/63, at night a person must be within 7 yds of the suspect to correctly identify a handheld object 100% of the time. If the officer's acuity is 20/100, the viewing distance reduces to 4.7 yds.


**Figure XI-1.** Percent of subjects (n=5) who were able to identify a handgun, cell phone, brown beer bottle and a pair of aviator-style black sunglasses in the daytime or purse at night with the various visual acuities and viewing distances during the day and night. (Carmean, et al., 2000)



In related research, Good and Augsburger (1987) decorrected 50 Columbus, Ohio patrol officers with 20/20 vision or better and asked them to identify whether a life-size target 20 ft (~7yds) away was holding a firearm. To simulate night conditions (when most shootings in Columbus were found to occur), the trials were conducted under low-light conditions (10 cd/m<sup>2</sup>). This resulted in a task that was moderately difficult even without decorrection. As indicated in Figure XI-2, without decorrection, officers misidentified 5-15% of the 60 targets presented. With vision between 20/30 - 20/40, the error rate increased to 15-25%. At 20/50 - 20/60, the error rate increased to 25-40%. An acuity of 20/45 was determined to be minimally acceptable, based on the convention that the threshold level of performance should fall halfway between the guess rate and zero errors. In this experiment, the guess rate was 50% because officers were given

two possible choices; therefore, 25% serves as the threshold level of performance. The resulting acuity for this error rate was 20/45.


**Figure XI-2.** Number of errors in identifying gun vs. no gun from a distance of 20 ft (~7 yds) under nighttime viewing conditions. Error bars are  $\pm$  one standard deviation. (Good & Augsburger, 1987)



Good et al. (1998) conducted a similar study based on a domestic disturbance. In this scenario, a person had either a firearm (long barrel shotgun, pistol-grip shotgun, handgun) or a distractor (umbrella, golf club, plastic spatula), and the officer had their spectacles dislodged in an altercation. The viewing distance was 20 feet, light levels were representative of indoor light levels, and viewing time was limited to two seconds. They also assessed an individual's ability to locate spectacles on a floor from a viewing distance of 8 ft in low light levels.

Figure XI-3 shows that weapon identification at acuities of 20/160 and lower were significantly worse than no-blur condition ( $p \leq .05$ ). Errors in finding spectacles began at 20/125 and approached chance levels at 20/200 for 80% of the participants. Based on the combined results, they concluded that an uncorrected visual acuity of 20/125 is an appropriate standard when consideration is given to spectacles becoming dislodged during a domestic disturbance call.

**Figure XI-3.** Mean correct responses in identifying whether a person has either a firearm (long barrel shotgun, pistol-grip shotgun, handgun) or a distractor (umbrella, golf club, plastic spatula) under typical room illumination levels. Error bars are  $\pm$  one standard deviation. (Good et al., 1998)




In a fifth study involving weapon identification, Johnson and Brintz (1996) decorrected six California Youth Authority supervisors and counselors whose vision was 20/20 or better. The simulation was conducted under night lighting (5 to 7 cd/m<sup>2</sup>) in an open dormitory setting. This luminance range was similar to the light levels used by Good and Augsburger (1987), but brighter than those used by Carmean et al. by a factor of at least 10. Fifteen surrogate wards were situated 5-7 ft. (2 yards) away from the participants. In each trial, one ward was holding either a weapon (knife or screwdriver) or a non-weapon (toothbrush or comb). The participants were tasked with detecting which ward was holding an object and identifying whether the object was a weapon or non-weapon.

With visual acuity at 20/20, there was 100% correct detection of which ward was holding an object (Figure XI-4). Detection fell to 80% correct for the 20/60 and 20/100 acuity levels, 60% at 20/200, and 20% at 20/400. The ability to identify whether the object was a weapon or non-weapon declined more rapidly with reductions in visual acuity. Correct identification at the 20/20 level was 75%, which degraded to 40% at 20/60, 25% at 20/100, less than 10% at 20/200, and 0% at 20/400.

Correct identification of only 75% with 20/20 vision from two yards indicates that this visual task was very demanding relative to the other weapon vs. no weapon studies. At this viewing distance, performance dropped when acuities were reduced to 20/60. The previous studies, particularly the Carmean et al. weapon identification experiment, would suggest that identification with 20/20 acuity at a 2-yard viewing distance should be nearly 100% correct, even at the much lower light levels used in the Carmean et al. study.

**Figure XI-4.** Average correct responses for object detection and weapons identification as a function of visual acuity. (Johnson and Brintz, 1996)



The difference in these results is most likely attributable to the difference in the choice of weapons. The knives and screwdrivers used in the Johnson and Brintz study were smaller than a handgun or shotgun and perhaps more similar in appearance to the non-weapons. Given that the task was near the limit of resolution for a person with 20/20 acuity, it is reasonable that a reduction in visual acuity to 20/60 made a large difference in performance, as compared to the smaller changes found in the other studies in which objects were sufficiently above the limit of resolution for those with 20/20 acuity.

The Johnson and Brintz study shows that although a person with 20/60 acuity may be able to detect a small object 80% of the time, an officer would have to be within 0.7 yards (arm's length) in order to identify it correctly 75% of the time.

The five studies together highlight the challenge of identifying a weapon at night, even for officers with 20/20 vision. In general, performance is significantly worse in night conditions at a 7-yard viewing distance when acuity falls below 20/40, due in large part to a reduction in the ability to resolve detail in low light levels. Johnson et al. (1992) found that 20/20 vision is degraded to 20/60 under typical night lighting conditions (i.e., sodium vapor streetlights), and 20/60 vision is degraded to 20/200 at night. Weapons identification improves in daylight or normal room illumination. In better lighting, a gun can be identified with reasonable accuracy from 7 yards with acuity near 20/100.

## 2. Facial Recognition

The recognition of a face or facial expression is another critical job skill. When pursuing or trying to recognize a suspect in a crowded area, only the suspect's face may be visible. Recognizing

and recalling facial features is also important when identifying a suspect in a line-up or when testifying in court.<sup>7</sup>

Most of the research into face and facial expression recognition has maintained a fixed distance and varied the contrast of the object, or has been concerned with computer algorithms that mimic human performance in recognizing faces. Three studies varied the angular size (i.e., viewing distance) and assessed the ability of individuals to recognize faces or their expressions.

First, Sheedy (1980) reported that facial recognition became difficult in low light levels when acuity became worse than 20/40. This result, however, was based on one subject (himself) and was carried out in lighting conditions that simulated urban street light levels.

Second, McCulloch et al. (2011) measured facial recognition as a function of optical blur using synthetic faces. The task was to identify which of four faces was different from the other three displayed at the same time. The faces were clearly unnatural in appearance; however, the rendering of a face into its basic components allowed the researchers to establish a feature difference that was quantifiable and set so that the difference between the target face and the three distractors was easily recognized in the no-blur condition. The face size was then scaled to simulate different viewing distances for the different levels of blur. Viewing conditions were comparable to a well-lit office.

The third study evaluated the ability of individuals with normal and reduced vision (due to macular degeneration) to recognize facial expressions at varying distances (Bullimore et al., 1991). Subjects were asked to identify one of four different facial expressions depicted by models in photographs in lighting conditions similar to a well-lit office.


Figure XI-5 displays the results of the McCulloch and Bullimore studies of viewing distances as a function of visual acuity for facial recognition. The error bars are at the 95% confidence interval for the McCulloch mean values. The mean threshold values are based on slightly different percent correct responses. The Bullimore thresholds are based on a 50% correct performance; the McCulloch thresholds are based on 63% correct. The results are very similar, despite slightly different threshold criteria, facial stimuli, and visual task (facial expression vs. facial recognition). This is consistent with Bullimore's conclusion that the visual acuity required for facial recognition and facial expression is similar as long as visual acuity loss is not large. The similarity of the findings for the two tasks indicates that the same data can be used to determine the visual acuity demands for both facial recognition and facial expression identification. The similarity of results is also remarkable given the different methods for reducing acuity (i.e., optical blur versus disruption of the central field due to age-related degeneration).

The equivalent viewing distances in Figure XI-5 indicate that a person with 20/40 acuity can recognize faces/expressions at a 7-yard (20 ft) viewing distance 50%- 63% of the time. Unfortunately, neither study indicated the values needed for 100% accuracy, although data from McCulloch et al. (2011) auxiliary experiments suggest that acuity values would need to be doubled to achieve a score of at least 90% correct. Therefore, an officer requires a visual acuity of 20/20 to recognize faces with 90% accuracy at a distance of 7 yards.

---

<sup>7</sup> As with the other visual tasks discussed above, facial recognition at a distance or in poor illumination can be affected by numerous factors in addition to visual ability per se; for example, race [whites have difficulty identifying black faces; blacks recognize white and black faces equally well (Cross, et al., 1971)], age [less errors with subjects of same age (Mason, 1986), and gender [less errors with subjects of same gender (Ellis, et al., 1973)].

**Figure XI-5.** Viewing distance as a function of visual acuity for facial recognition (McCulloch et al., 2011) and facial expression recognition (Bullimore et al., 1991). The error bars are the 95% confidence intervals for McCulloch et al.'s results and the dashed curve is the average of the two studies' results.



### 3. License Plate Identification

The ability to read and identify license plate numbers from a distance is another essential job function. When in pursuit of a vehicle at 60 mph, maintaining a safe distance (i.e., 6 car lengths) requires that the officer read the plate from a distance of 100 ft. Sheedy (1980) observed that he could read a license plate from this distance with 20/20 vision and good lighting conditions. By extrapolation, someone with 20/40 acuity would be unable to read a license plate from 50 ft (3 car lengths) away. Sheedy noted that these distances assume no movement; under dynamic conditions, viewing distances would be even shorter. He also pointed out that his sighting distance was shorter than predicted by a factor of 1.53, based on the dimensions of the numbers and his acuity of 20/20. He attributed the difference to the narrow width and lower contrast of the license plate numbers and letters.

Given the difficulty of judging car lengths, a 2-3 second interval between cars passing roadway landmarks is commonly adopted as a safe following distance. Assuming a suspect's car and police cruiser are traveling at 55 mph and the license plate characters are 2.75 inches high, the separation time for an officer with 20/20 acuity would need to be 1.96 sec (158 ft) or less to read the rear license plate characters in daylight. If the license plate characters were lower in contrast, such as those viewed by Sheedy, the separation time would be reduced to 1.3 sec (105 ft). If the acuity was 20/40, the separation time would be further reduced to 0.98 - 0.65 seconds, depending on the plate characteristics. The task is even more challenging at night: the acuity of an officer with 20/20 acuity in daylight conditions would be reduced to 20/25 under conditions similar to reading license plates at night (Hovis & Ramaswamy, 2006). The resulting separation time would be reduced to 1.04-1.57 seconds, depending upon the characters on the plate.

### 4. Withdrawal Under Fire

An officer must have sufficient vision to withdraw from a dangerous situation and seek safety. In a study conducted by the Canadian military (Gibbs, 2001), soldiers had to withdraw from trenches to armored personnel carrier. The time required to withdraw increased significantly when the acuities were reduced to 20/200 or 20/400; however, the withdrawal times for 20/20

and 20/120 were similar. The relative increase in time for the two lower acuities was 9% (10 sec). These data suggest that seeking safety could be compromised when the acuity level drops below 20/120.

Table XI-4 summarizes the results of the above-referenced decorrection studies on the impact of visual acuity on critical peace officer task performance. As indicated in that table, unimpaired visual acuity is necessary for quick identification of objects at varying distances. **A 20/20 vision standard is therefore justified, given an officer's need for facial recognition, firing weapons at distant targets, and driving.** The need for unimpaired vision is even more compelling for officers who may be called upon to perform these duties at night.

**Table XI-4. Critical Task Performance Determined by Decorrection Studies**

| VISUAL ACUITY | CRITICAL TASK PERFORMANCE  |
|---------------|--|
| 20/20         | <ul style="list-style-type: none"> <li>● In good light, can consistently identify weapons at distances of up to 25 yards<sup>1</sup></li> <li>● In low light, will identify guns correctly at 7 yards with an error rate of 0-15%<sup>2,6,7</sup></li> <li>● Under night conditions, from 5-7 feet can detect whether an individual is holding a small object with 100% accuracy and can identify an object (comb vs. knife) with 75% accuracy<sup>3</sup></li> <li>● Facial identification with 75% accuracy at 14 yards<sup>4,8</sup></li> <li>● License plate identification from 100 feet (6 car lengths)<sup>5</sup> to 150 feet</li> </ul>   |
| 20/30         | <ul style="list-style-type: none"> <li>● Facial identification with 50% to 63% accuracy at 9 yards<sup>4,8</sup></li> </ul>  |
| 20/40         | <ul style="list-style-type: none"> <li>● In good light, can consistently identify guns at 7 yards;<sup>1,6</sup> at 15 yards the error rate increases to 14%<sup>1</sup></li> <li>● In low light, can identify guns at 7 yards, but the error rate can be as high as 25%<sup>2,7</sup></li> <li>● Legal limit for driving any vehicle without further assessment of visual capabilities</li> <li>● License plate identification from 50 feet (3 car lengths)<sup>5</sup> to 75 feet</li> <li>● Facial identification: accuracy is 50% - 63% at 6 yards<sup>4,8</sup></li> </ul>  |
| 20/50         | <ul style="list-style-type: none"> <li>● In low light, can misidentify guns at 7 yards with an average error rate of &gt;25%<sup>2</sup></li> <li>● Facial identification accuracy is 50% to 63% at 5 yards<sup>4,8</sup></li> </ul>   |
| 20/60         | <ul style="list-style-type: none"> <li>● Under night conditions, from 2 yards can detect whether an individual is holding a small object with 80% accuracy, but can only identify the object (comb vs. knife) with 40% accuracy<sup>3</sup></li> <li>● Under night conditions, errors in identifying a gun can be &gt;25%<sup>2</sup></li> <li>● Facial identification is 63% accuracy at 5 yards<sup>4,8</sup></li> </ul>   |
| 20/80         | <ul style="list-style-type: none"> <li>● In good light, can identify guns at 7 yards with an error rate of 8%; 22% is the error rate at 15 yards<sup>1,6</sup></li> <li>● In low light, will misidentify guns at 7 yards with an average error rate that is &gt;30%<sup>2</sup></li> <li>● Facial identification possible with 50% to 63% accuracy only at 3 yards<sup>4,8</sup></li> <li>● License plate identification at 25 to 38 feet<sup>5</sup></li> </ul>   |
| 20/100        | <ul style="list-style-type: none"> <li>● Under night conditions, from 2 yards, can detect whether an individual is holding a small object with 80% accuracy and can identify an object (comb vs knife) with 25% accuracy<sup>3</sup></li> <li>● Identification of guns from 7 yards is impaired in both good and poor lighting<sup>1,2,6,7</sup></li> </ul>  |
| 20/200        | <ul style="list-style-type: none"> <li>● In good light, can identify guns at 7 yards with error rate between 17% and 33%;<sup>1,6</sup> 39% error rate at 15 yards<sup>1</sup></li> <li>● In low light, identifying guns at 7 yards approaches the guess rate<sup>2,7</sup></li> <li>● Under night conditions, from 2 yards, can detect whether an individual is holding a small object with 60% accuracy but can identify the object with less than 10% accuracy<sup>3</sup></li> <li>● Facial identification is impossible beyond an arm's length<sup>4</sup></li> <li>● License plate identification impossible at &gt;10 feet<sup>5</sup></li> <li>● Locating spectacles from 2.5 yards is by chance<sup>6</sup></li> <li>● Legal blindness as defined by the Social Security Administration and the IRS</li> <li>● Withdraw under fire in the daytime is compromised</li> </ul> |

<sup>1</sup>Giannoni, 1981; <sup>2</sup>Good & Augsburger, 1987; <sup>3</sup>Johnson & Brintz, 1993; <sup>4</sup>Bullimore et al., 1991; <sup>5</sup>Sheedy, 1980; <sup>6</sup> Good et al., 1998; <sup>7</sup>Carmean et al., 2000; <sup>8</sup>McCulloch et al., 2011; <sup>9</sup> Gibbs, 2001



## b. FAR ACUITY STANDARDS FOR EACH EYE VS. BOTH EYES


In order to justify an "each eye" standard, it must be shown that poor acuity in the weaker eye could hamper the safe performance of peace officer functions. Three job demands and circumstances that would influence a standard for each eye are: (1) use of a shotgun or long gun; (2) peripheral vision; and (3) trauma to one eye that results in sudden loss of vision.

### 1. Use of a Shotgun or Long Gun

Certain tactical situations may require the use of a shotgun or long gun from behind either a left-handed or right-handed barrier (such as a doorway). Figure XI-6(A) illustrates the proper stance. If the officer's vision is impaired in the sighting eye (in this illustration, his left eye), it would be very difficult to properly aim, resulting in degraded accuracy. The alternative would be for the officer to sight with the better eye and shoot with the opposite hand [Figure XI-6(B)]. This positioning would require that the officer expose most of the body and the entire head, creating a safety risk.

For agencies that use long guns or shotguns, an acuity of at least 20/40 in the weaker eye is necessary to allow for adequate identification of weapons and facial recognition within 7 yards under most lighting conditions.

**Figure XI-6.** Use of a long gun behind a left barricade. (A) The correct tactical stance requires aiming the weapon with left eye. (B) Aiming weapon with right eye exposes officer to fire. (Photos courtesy of the Training Division, Los Angeles Police Department)



### 2. Peripheral Vision

Peripheral vision requires functional vision in each eye. Peripheral vision is discussed under [Visual Field Deficiency](#).

### **3. Trauma to one eye with sudden loss of vision**

The likelihood of losing vision in one eye during a critical incident due to sudden trauma also justifies a minimum far acuity requirement for each eye. In a 1984 survey of 158 California police officers, 41.2% of officers reported an eye injury while on patrol. Based on the average length of service, the annual rate of eye injuries was estimated to be 25 per 300 officers. If exposure to noxious chemicals is excluded (since this is likely to impair vision in both eyes), the annual rate is reduced to 16 incidents per 300 officers (Briggs, 1984).

The rates of serious eye injuries (i.e., those that result in compensation claims) are lower. LAPD worker's compensation records for the years 1987-1990 revealed that unilateral eye injuries during altercations occurred at an annual rate of approximately 1 per 300 officers assigned to field duty (Goldberg, 1993). The annual rate of all eye injuries for Milwaukee police officers was 4.5 per 300 officers during 1992-1993, and approximately 1 per 300 officers between 1996- 2008 (Brandl, 1996; Brandl & Stroshine, 2012). Eye injuries resulting specifically from assault or suspect-related events were 1.4 per 300 officers, but this only reflects a one year (1992-1993) time frame (Brandl, 1996).

The relatively low likelihoods associated with a traumatic loss of vision in one eye must be balanced against the significant consequences of such an event. Although the incidence of self-reported eye injuries is significantly higher than those injuries that resulted in workers' compensation claims, it cannot be surmised that these self-reported injuries were not serious. It is reasonable to assume that the officer's vision in one eye was either partially or temporally affected, requiring reliance on the other eye during the incident.

Visual acuity of 20/20 for the better eye and 20/40 for the weaker eye should be required of officers whose responsibilities include the use of long guns or shotguns, or that require activities where they may need to rely on either eye separately. For officers who do not use these weapons, the acuity requirement for the weaker eye should be 20/125. This requirement would allow the officer to retreat quickly, detect the presence of small hand-held objects at a viewing distance of two yards under low light levels (Johnson & Brintz, 1993), and identify the presence of guns and knives from a viewing distance of 4.7 yds with over 80% accuracy in low and bright light levels (Giannoni, 1981; Carmean et al., 2000; Good et al., 1998).

#### **c. METHODS OF CORRECTION**

The uncorrected vision of a significant proportion of the population falls short of 20/20. Among a sample of 200 LAPD applicants, for example, 32% were found to have uncorrected vision of less than 20/20; even a far acuity standard of 20/30 uncorrected would eliminate 19% of this sample (Table XI-5).

Various methods exist for correcting vision, including spectacles, contact lenses, orthokeratology, and refractive surgery. Each method has its attendant advantages and risks. This section discusses factors to consider when determining the acceptability of each method.

**Table XI-5:** Distribution of Uncorrected Vision in 200 LAPD Applicants with Both Eyes Open

| Uncorrected Vision* | Percent of Applicants With This Level of Vision or Better |
|---------------------|---|
| 20/20               | 68%   |
| 20/25               | 75%   |
| 20/30               | 81%   |
| 20/40               | 83%   |
| 20/50               | 86%   |
| 20/80               | 90%   |
| 20/200              | 94%   |

\*Single character errors were ignored except at the 20/200 level; 20/40-1 was considered 20/40, 20/200-1 was considered to be worse than 20/200. Goldberg, R.L. (1993). Uncorrected vision of LAPD applicants. Unpublished data.

## 1. Spectacles

The level of risk posed by spectacle-wearing officers depends upon the probability that an officer would lose the use of his/her spectacles during a critical incident and the likelihood that this would result in impairment and/or injury. These concerns, in turn, must be balanced against the potential benefits, such as protection against thrown objects, sand, etc. Each of these issues is discussed below.

### ***What is the probability of an officer losing the use of spectacles while on duty, particularly during a critical incident?***

Spectacles can become dislodged and/or broken when an officer is assaulted by a resisting suspect, when pursuing a suspect, or when an officer is required to make a sudden vehicle stop. Climatic factors such as rain, snow, and lens fogging may also suddenly deprive an officer of full visual correction.

Results from surveys on spectacle wear and policing bear significant similarities. In a survey of 195 myopic LAPD officers, Mancuso (1987) found that eighty-six officers (44%) indicated that they had been involved in an incident where they needed to see without their spectacles (Table XI-6). Approximately 28% of the officers stated that this occurs less than once per year, 45% stated 1-6 times per year, 13% stated 7-20 times per year, and 14% stated more than 20 times per year. For the entire group (N=195), on average, each officer was required to function without glasses approximately twice per year during a critical incident.

A similar questionnaire survey was conducted on 292 officers from the City of Columbus, Ohio (Good and Augsburger, 1987). Fifty-two percent of the officers reported that their glasses dislodged during the performance of their duties at least once in their career, yielding a probability of dislodgement of 34% per year per officer. The Ohio survey also found that 67% of officers reported that they have had to remove their glasses because of rain or snow at least once in their career; 56% reported removing their glasses due to fogging.

**Table XI-6: 1987 LAPD Vision Questionnaire of Incumbent Police Officers**

|   | <b>% Answering Yes<br/>(Spectacles)</b> |
|---|---|
| Have you ever sustained an on-the-job injury specifically related to your wearing your corrective lenses?   | 5%<br>(10/194)                          |
| Have you ever been involved in critical incidents, including but not limited to the apprehension of suspects, physical altercations, or vehicle pursuits, which necessitated that you see without your corrective lenses? | 43%<br>(83/195)                         |
| Has your wearing corrective lenses ever been an issue during a court appearance?  | 15%<br>(27/184)                         |
| Do you believe that wearing corrective lenses presents an imminent hazard to your safety, that of your co-workers, or that of the public in any way?  | 6%<br>(12/195)                          |
| Have you ever encountered any job safety problems caused by your corrective lenses?   | 28.9%<br>(57/197)                       |

Mancuso, 1987. Responses of myopic LAPD officers to a vision questionnaire. Unpublished study.

These findings and those of three other peace officer surveys on spectacle removal while on duty, are summarized in Table XI-7. As indicated in that table, the percentages of officers reporting at least one dislodgement are very similar. There is considerable variance in the percentages of officers removing their spectacles because environmental factors (lens fogging, snow and rain); it is not surprising that RCMP officers reported the highest percentage of removing spectacles due to environmental factors throughout Canada. The other result of note is that the percentages within each row sum to values greater than 100% for most of the studies, indicating that officers have their spectacles removed/dislodged several times in their career for different reasons.

**Table XI-7. Percentage of spectacle wearers responding “yes” to questions about spectacle wear while on duty.**

| <b>Study,<br/>Location,<br/>Number of spectacle<br/>wearers</b> | <b>Spectacles<br/>dislodged at<br/>least once<br/>during career</b> | <b>Spectacles<br/>broken at least<br/>once during<br/>career</b> | <b>Spectacles had<br/>to be removed<br/>because of<br/>environmental<br/>factors</b> |
|---|---|--|--|
| Good and Augsburg, 1987<br>Columbus OH<br>N=292                 | 52%   | N/A  | 61%  |
| Hovis et al., 1998<br>Ontario Canada<br>N=66                    | 53%   | 37%  | 51%  |
| Wells et al., 1997<br>RCMP<br>N=574                             | 58%   | 46%  | 76%  |
| Good et al., 1998<br>Columbus OH<br>N=403                       | 49%   | N/A  | 22%  |

There have been two published studies of spectacles reimbursement rates. Sheedy (1980) reported that during a two-year period, the City of Columbus, Ohio, reimbursed eight officers for spectacles broken during altercations. Giannoni (1981) reported that during fiscal year 1979-80, the CHP reimbursed 17 officers for spectacles broken during altercations and two officers who lost their spectacles during foot pursuits (Table XI-8). Unfortunately, neither study provided data

on the total number of spectacles-wearing officers, so calculations of the relative rates of loss or breakage are not possible.

**Table XI-8.** Number of prescription eyeglass reimbursement requests submitted by CHP during 1979-80 by job-related loss or breakage categories

| Category                                | Number of Reimbursement Requests |
|---|----------------------------------|
| 1. Assault/resisting arrest             | 17                               |
| 2. CHP patrol car/motorcycle accident   | 4                                |
| 3. Removing debris on highways/freeways | 1                                |
| 4. Accident investigations              | 3                                |
| 5. Rescue/first aid                     | 4                                |
| 6. Foot pursuits                        | 2                                |
| 7. Operating motorcycle                 | 2                                |
| 8. Routine stop                         | 5                                |
| 9. Other*                               | 9                                |

\*Fall on pavement, sparks from battery, etc.

(Giannoni, Entry-level vision requirements validation study. Personnel Bureau, California Highway Patrol. October 1981)

Dodson (1993) and others have argued that the risk of officers losing their spectacles can be virtually eliminated by the use of military spectacles and other devices that secure spectacles to the head. Unfortunately, retaining devices such as straps and cords can be a potential safety hazard since, during an altercation, they could be used to choke the officer. Also, spectacles held tightly by elastic, as is with athletic eyewear, could be forcibly snapped back into the officer's face. Moreover, it is unlikely that the tight elastic would be tolerated for an 8-hour shift.

Newer types of combat frames that are secured by a "D" shaped earring are uncomfortable when fitted tightly enough to avoid dislodgement during altercations -- a light tapping to the side of the frame can cause severe pain to the bridge of the nose. Although more attractive than traditional military frames, the newer generation of combat spectacles are very conspicuous and relatively unattractive, which could hamper their acceptance, use, and public reaction.

**Note:** All spectacles worn by officers on duty should consist of polycarbonate lenses and frames that meet ANSI Z87.1 specifications. This greatly reduces the likelihood and severity of injury to the officer.

***How often would the loss of spectacles result in injury or other negative consequences?***

It has been argued that losing one's spectacles during a critical incident would be unlikely to result in negative consequences for all but the severely myopic, since a suspect is usually situated very close to the officer in these situations (Holden, 1993; Dodson, 1993). Situations such as these may be further mitigated by the presence of a partner and/or the potential availability of a spare pair of spectacles. However, a study conducted for the California Youth Authority showed that refractive error affects the visual detection and identification of weapons even at distances as short as 5-7 ft (Johnson & Brintz, 1993). Even those who advocate for this position acknowledge the seriousness of the consequences that could (and do) occur in these situations. Holden (1993) reports an incident in which the loss of spectacles is believed to have contributed to the death of an FBI agent.

Dodson (1993) recommended that myopic officers wear combat spectacles and be provided with handguns that have special high-visibility sights; however, approximately 70% of the time,

officers do not use sight alignment (Sheedy, 1980). In addition, standard clinic practice for professional and sport shooters is to provide a correction so that gun's far sight is in best focus, which would be unnecessary for most nearsighted individuals, as their location of clear focus is approximately arm's length.

In a 1984 POST survey, 53 spectacles-wearing officers from various agencies were asked to report any negative experiences (including but not limited to impairment or personal injury) associated with wearing spectacles while on duty. As indicated in Table XI-9, the percent of negative outcomes reported by spectacle wearers was less than 4%. The majority of physical harm outcomes were cuts and bruises to the officer, and the majority of property damage outcomes consisted of broken spectacles. All dislodgements occurred during an arrest or altercation with a suspect. This is equivalent to an annual risk per officer of approximately 1.1% (average length of service = 5 years).

The POST survey also asked a larger group of officers whether they knew of other officers who experienced the same array of negative consequences on the job due to the use of spectacles. Table XI-9 reports these results for 140 officers. Although such questions generated a large number of anecdotal cases (rather than incident rates), the percentage witnessed by officers is similar to the self-reported rates. The one major difference between the two survey groups was that two officers witnessed an automobile accident in which the dislodgement of spectacles could have been a contributing factor.

Sheedy (1980) reported that during a two-year period, there was one report where an officer's spectacles were broken, and because of his poor vision, the suspect escaped. If it can be assumed that there are 109 spectacle wearers (based on Good and Augsburger's 1987 survey of the same police force approximately 10 years later), the estimated annual incident rate is 0.5%.

**Table XI-9:** Reported Instances of Negative Consequences Resulting from the Use of Spectacles by Officers

| Impairment | % Self-report (N=53) | % Witnessed (N=140) | Outcome  | Circumstances   |
|------------|----------------------|---------------------|--|---|
| Chemicals  | 1.9%<br>(1)          | 0                   | Failure to provide required duty (make an arrest, complete search) | Maced in combative situations--arrest delayed   |
| Dislodged  | 0                    | 1.4%<br>(2)         | Failure to provide required duty (make an arrest, complete search) | Spectacles dislodged while making an arrest   |
| Dislodged  | 3.8%<br>(2)          | 4.3%<br>(6)         | Physical harm to self or others                                    | Altercation – officer struck in face; spectacles broken, knocked off, cut officer's face; suspect had to be subdued by other officers |
| Dislodged  | 3.8%<br>(2)          | 5%<br>(7)           | Loss or damage of property   | Resulted in automobile accident, broken spectacles  |
| Fogged     | 1.9%<br>(1)          | 1.4%<br>(2)         | Failure to provide or delay in service                             | Cold to warm; had to clean; other officers had to assist; fogged unable to see  |

Two later surveys asked spectacle-wearing officers if they felt that their safety or that of the public was compromised when their spectacles were dislodged or removed (Hovis et al., 1998; Good et al., 1998). The percentage that responded positively to these questions ranged from

5%-13%. In addition, 14%-22% of the spectacle wearers in the Good survey reported that their efficiency was compromised without their spectacles. The exact annual incident rate for the spectacle wearers could not be calculated from either study, but the Ontario value was estimated to be 0.6% (Hovis et al., 1998).

These results do indicate safety concerns when officers have their spectacles dislodged or removed while on duty. The potential for a negative outcome appears to be near 1% per year for spectacle wearers. However, it must be remembered that the agencies had uncorrected visual acuity requirements at the time of this research. For example, the Columbus police required uncorrected acuity of 20/40 (although some officers were hired under an earlier 20/50 standard). The uncorrected acuity standard for the Ontario police agencies varied between 20/40 - 20/60.

The degree of risk associated with wearing spectacles is directly proportional to the candidate's degree of visual impairment (see Table XI-4); therefore, it is reasonable to conclude that spectacles are acceptable for candidates with relatively mild degrees of uncorrected refractive errors.

### ***How often do spectacles provide protection from hazards?***

The POST survey also studied the potential benefit of spectacle wear. The 53 officers who wore spectacles listed over 50 incidents in which they felt that spectacles protected them from injury (Table XI-10). Some of these incidents involved confrontations with suspects who tried to disable the officer by throwing sand or other matter into the officer's face. The results from officers in the study who did not wear spectacles were similar. Incidents where the spectacles provided protection from sand, dust, gravel and rocks, whether from the natural environment or thrown by suspects, were so numerous that the number of incidents was not tabulated.

**Table XI-10:** Reported Instances Where Spectacles Provided Officers Protection

| <b># Times</b> | <b>Circumstances</b>   |
|----------------|--|
| 1              | Broken windshield -- eyes protected from glass   |
| 5              | (1) Lead splatter at range<br>(2) Wall particles -- removing evidence<br>(3) Dura print fumes                                  |
| 4              | (1) Flying objects<br>(2) Leaking chemicals in a fire  |
| 5              | Tear gas, objects thrown, struck in face, spit on  |
| 1              | Suspect threw sand -- spectacles protected eyes  |
| 5              | Spectacles protected eyes from thrown gravel   |
| 10             | Spectacles acted as shield for eyes  |
| 10             | Prevented dust or hard objects from entering or harming my eyes  |
| Many           | Objects thrown, i.e., dirt, sand, etc., by people and natural forces. Also limbs, branches, bushes scratched face but not eyes |
| -              | Strong winds -- debris hit spectacles  |
| Several        | Protection from windblown dust/dirt  |
| 3              | Blowing sand in two storms. Blowback from weapon on range  |
| 4              | Protection against blowing sand/debris from helicopter blade thrust  |
| Many           | Sand/rocks/bugs while a motorcycle officer   |

Briggs (1984). POST Study of Vision-Related Characteristics/Experiences. Unpublished report.

**SUMMARY:** An uncorrected visual standard of 20/40 in each eye is recommended for spectacles-wearing officers for officers required to use long guns or shotguns. For all officers, serious eye injuries, although infrequent, occur as a result of an assault in 1/3 of cases, and it is uncertain as to whether the officers were performing critical tasks when the other 2/3 of the injuries occurred. Therefore, an uncorrected standard of 20/40 in the better eye and between 20/40 - 20/125 in the worse eye should be established. A visual acuity worse than 20/125 would leave an officer markedly impaired without spectacles (Table XI-4).

## 2. Contact Lenses

Contact lenses can be classified by their rigidity:

**Soft Contact Lenses (SCL)** are large, flexible and permeable to oxygen. Advantages include better comfort, low risk of dislodgement, low risk of particle entrapment, and extended wear availability. Disadvantages include the need for regular cleaning/disinfection if the SCLs are not daily or extended wear disposable lenses. SCLs are worn by approximately 90% of contact lens wearers (Nichols & Starcher, 2020)

**Rigid Gas Permeable Lenses (RGP)** are made of a rigid material that is permeable to oxygen (Key, 1990). There are two types of RGP lenses: (1) the *corneal RGP* lens (corneal RGP) is smaller in diameter (~9 mm) than the candidate's visible iris; (2) the *scleral RGP* lens (scleral lens) (diameter varies from 14 mm to 20 mm) vaults the cornea and rests on the scleral conjunctiva (Bowden & Barnett, 2017). The advantages of both types of lenses include easier care (no disinfection required) and the ability to correct for corneal irregularities (Schornack, 2017). Disadvantages of the corneal RGP lens include discomfort, easy dislodgement, and a high risk of particle entrapment. Discomfort, particle entrapment, and potential for dislodgment are less of a concern with scleral lenses since they are stable, cover the entire cornea and land on the scleral conjunctiva.

**Hybrid Contact Lenses (HCL)** are composed of a gas permeable center and soft skirt with high oxygen transmission. The newest generation of these lenses has overcome problems of durability and irritation of the cornea and conjunctiva (Maguen, et al., 1991,1992). However, it is possible that the soft skirt could separate from the RGP center in approximately 5% of the wearers (Harbiyeli, et al., 2020). Like the scleral lens, the HCL can correct corneal irregularities, and the potential for dislodgment compared to corneal RGPs is less of a concern with HCLs since they are stable and cover the entire cornea. Disadvantages include the need for regular cleaning/disinfection.

Both scleral lenses and HCLs can be used to correct all types of refractive errors, including large amounts of regular astigmatism (Piñero et al., 2015; Abou Samra, et al, 2018; Barnett et al., 2018). However, the current practice is to prescribe scleral lenses and HCLs in cases where there are corneal irregularities and dry eye disease (Nau, 2008; Schornack , 2017; Sarac et al., 2019; Schornack, et al., 2019). Since some of these conditions are progressive, additional questions regarding the stability of the refractive error, corrected visual acuities and uncorrected visual acuities are warranted.



Two issues must be considered when determining the acceptability of contact lens use by peace officers: safety and compliance.

a. **Safety.** The use of contact lenses could potentially create a safety hazard under the following circumstances:

- (1) **Lenses lost during an altercation.** Table XI-11 displays the results of six studies that researched the percentages of SCL and corneal RGP contact lens-wearing police officers who reported lens removal, dislodgement or fogging while on duty. The results indicate that contact lens dislodgements are less frequent than spectacle dislodgments. Dislodgement of both lenses is very rare: in a survey of 336 RCMP officers who wore contact lenses, only 1.4% reported that it occurred during their career (Wells et al. 1998). However, in general, the percentages of officers reporting that they removed their lenses due to irritation approaches that of spectacle wearers.

In the Mancuso study, four officers (10%) reported occurrences in which they were not wearing contact lenses during a critical incident. One officer had this happen less than once per year; another officer reported occurrences of 1-6 times per year. The remaining two officers reported occurrences of more than 6 times per year.

**Table XI-11: On-Duty Loss of Use of Contact Lenses**

| Study, Location, Number of Contact Lens Wearers   | Contact lenses dislodged at least once during career                      | Contact lenses blurred vision due to fogging/ freezing or other factors | Contact lenses had to be removed because of environmental factors | Lost a contact lens while on duty        |
|---|---|---|---|--|
| Briggs, 1984<br>N=17                              | 8% (soft)<br>0% (hard and corneal RGP)                                    |   |   |  |
| Mancuso, 1987<br>LAPD<br>N=38                     | 11% (all lens types)<br>(Perform without corrective lenses at least once) |   |   |  |
| Good and Augsburger, 1987<br>Columbus OH<br>N=108 | 19% (soft)<br>20% (hard and corneal RGP)                                  | 47% (soft)<br>60% (hard and corneal RGP)                                | 46% (soft)<br>57% (hard and corneal RGP)                          | 10% (soft)<br>14% (hard and corneal RGP) |
| Hovis et al., 1998<br>Ontario Canada<br>N=20      | 6% (soft)<br>5% (hard and corneal RGP)                                    |   | 50% (soft)<br>51% (hard and corneal RGP)                          |  |
| Wells et al., 1998<br>RCMP<br>N=330               | 21% (all lens types)  | 30% (all lens types)  | 37% (all lens types)  | 9.8% (all lens types)                    |
| Good et al., 1998<br>Columbus OH<br>N=183         | 31% (all lens types)  |   | 52% (all lens types)  | 16% (all lens types)                     |

Because the most recent designs of scleral lenses and HCLs are relatively new, similar data for these lenses in the policing environment is not available. Based on the fact that scleral lenses and HCLs are similar in size to the SCL and more stable than corneal RGPs, the potential for particle entrapment and dislodgement for these two lens types should be similar to the SCLs.

Nevertheless, scleral lenses may be prone to midday fogging and the wearer may have to remove the lenses for cleaning and refill the fluid reservoir. Between 7% and 49% of the wearers have to remove their scleral lenses at least once during the day, with 20% to 30% of the wearers being the most commonly reported values (Visser et al., 2007; Walker et al., 2015; Schornack et al., 2019; Fuller & Wang, 2020). Because midday fogging is relatively common, a majority of practitioners who fit scleral lenses recommend breaks during the day to remove, clean and reapply lenses at least some of the time (Harthan et al., 2018). As improvements in lens design and solutions evolve, the prevalence of midday fogging may decrease (Barnett et al., 2020; Fogt et al., 2020).

In general, reported safety concerns are lower than those for spectacle wearers. Nevertheless, contact lenses are still dislodged and/or have to be removed while on duty. The scleral lenses are susceptible to midday fogging.

- (2) **Use of contacts in hazardous environments.** Based on studies researching the use of soft contact lenses in hazardous environments, these lenses could be worn safely while carrying out policing duties (Kok-van Aalphen, 1985; Royall, 1977; Nilsson et al., 1981; Nilsson & Andersson, 1982; Rengstorff & Black, 1974).
- (3) **Particle entrapment under a lens.** Particle entrapment can result in acutely painful and incapacitating "contact lens attack." Vision in the nonaffected eye is markedly impaired due to sympathetic tearing and photophobia until the other lens is removed. Particle entrapment is more likely to occur with smaller diameter corneal RGP lenses. Surveys of police officers reveal that, due to irritation, RGP lenses are removed at higher frequencies than larger diameter SCLs (Hovis et al., 1998). The increased probability of particle entrapment for corneal RGP lenses has led the American Optometric Association to recommend against the use of these lenses in industrial environments (AOA, 1990); their use in military aviation is also discouraged due to the high levels of particulate in cabin air (Polse et al., 1990).

The available evidence suggests that soft, hybrid and scleral contact lenses can be used by peace officers with minimal risks. Soft, hybrid and scleral lenses are preferable to corneal RGP lenses due to a lower likelihood of particle entrapment. The resurgence of scleral and hybrid lenses made of RGP materials may offer a solution to the particle entrapment issue for those individuals who require or prefer RGP lenses.

## **b. Compliance**

- (1) **Temporary discontinuation due to eye infections, corneal abrasions, allergies, or other medical conditions.** Between 17% - 25% of contact lens-wearing peace officers (both corneal RGP and SCL wearers) reported that they could not wear their lenses for at least one day per year due to irritation, discomfort or temporary medical conditions. The median number of days ranged from 2-3 days per year (Hovis et al., 1998; Wells et al., 1998), with 2% of officers discontinuing contact lens wear for more than 50 days. This is comparable to the mean value of three days reported for the general population of SCL wearers (Nilsson & Lindh, 1984).

For persons who have worn soft contact lenses successfully for more than a year, motivational factors are more responsible for episodes of temporary discontinuation than medical complications. Between 1988-1993, the LAPD hired over 300 officers who had worn SCLs successfully for at least one year and who signed a pre-placement agreement (see Figure XI-7), obligating them to wear SCLs whenever assigned to field duty. During five random department-wide inspections conducted between June 1990

and November 1991, non-compliance rates varied between 2-8% on a given day, with an average rate of 5%. Thirty officers were found on duty without their SCLs on 39 occasions; five officers were non-compliant twice, and two officers were found non-compliant three times.

Medical reasons were cited for non-compliance in only six (15%) of the incidents. More commonly, officers said they forgot their contacts, lost one, or preferred to wear spectacles. There was a slight, nonsignificant increase in non-compliance in officers who had been on the job for longer periods of time (Figure XI-8). Discipline was limited to written reprimands and quarterly eye inspections had not been conducted regularly; presumably, non-compliance among these patrol officers could have been significantly reduced with stronger administrative controls.

Similar data for scleral lenses and HCLs in the policing environment is unavailable. Nevertheless, data from the general population suggest that temporary discontinuation of scleral lenses due to medical conditions would be under 9% (Pullum, et al., 2005; Walker et al., 2016; Fernández-Velázquez;2019; Fuller & Wang, 2020), although it could be as high as 18% depending on how the practitioner would choose to manage the adverse event (Schornack et al., 2019). Nevertheless, the incidence of temporary discontinuation of scleral lenses does not appear to be higher than the rates for SCL. The temporary discontinuation of HCL is harder to estimate because of the relatively small numbers of patients fit with these lenses; however, the available data suggests that the rate would be approximately 10% of the wearers (Harbiyeli et al., 2020), similar to that of SCL wearers.

- (2) ***Permanent Discontinuation.*** Discontinuation rates appear to be highest during the first year of use. Sulley and colleagues (2017, 2018) reported that approximately 25% of new SCL wearers discontinued within the first year. The most common reason (41%) for discontinuation was poor vision (distance and near) followed by discomfort (36%) and handling difficulties (25%). The discontinuation rates for the general population were higher compared with data on SCL-wearing USAF aircrew. Fifteen percent (15%) of the aircrew reported discontinuing lens wear in their first year. Similar to Sulley et al.'s (2018) results, unacceptable vision was the most common reason for discontinuing lens wear, accounting for 73% of the aircrew who stopped wearing lenses during the first year (Dennis et al., 1992).

Scleral lens wearers have similar rates of permanent discontinuation within the first few months of wear. The percentage of wearers who discontinue wear within the first 3 to 6 months is approximately 26% (Pullum et al., 2005; Pecego et al., 2012), although in a population with other comorbidities, the rate could be approximately 35% (Kanakamedala et al., 2019). The permanent discontinuation rate for HCL is lower at approximately 15% of the new fits (Abdalla et al., 2010; Abou Samra et al., 2018; Harbiyeli et al., 2020; Uçakhan & Yeşiltaş, 2020). Although these lenses are often worn by individuals with compromised corneae and anterior ocular surfaces, the discontinuation rates during the first few months of wear are similar to SCL wearers. As with SCL wearers, inadequate vision, comfort and handling issues were the primary reasons for discontinuing wear (Pecego et al., 2012; Kanakamedala et al., 2019; Abdalla et al., 2010; Abou Samra et al., 2018; Harbiyeli et al., 2020; Uçakhan & Yeşiltaş, 2020).

Figure XI-7. Sample Pre-Placement Agreement Involving Use of Contact Lenses

# SAMPLE

## PRE-EMPLOYMENT NOTICE OF CONTACT LENS REQUIREMENTS

Name: \_\_\_\_\_ Date of Hire: \_\_\_\_\_

**Medical Condition:** EXAMPLE: *Poor uncorrected distance vision - myopia correctable with soft contact lenses.*

*I acknowledge that the medical condition noted above was present at the time that the **(name of law enforcement agency)** offered me employment. I affirm that I am currently, and have been for the past six (6) months prior to employment, a bona fide, successful **(soft / scleral / hybrid)** contact lens wearer. I also understand that my use of these contact lenses is permitted as a reasonable accommodation for my **(medical condition)**.*

*I understand that my ability to perform the duties assigned to me as a full-duty patrol officer may be contingent upon my ability to wear successfully contact lenses on duty, and I shall wear such lenses whenever I am on duty except when authorized by my supervisor (or the Employee Assistance Unit) to do otherwise. I also understand that it is my responsibility to notify my supervisor (or the Employee Assistance Unit) should I become unable to wear contact lenses while on full duty or should I take any other medical action, which would otherwise affect my vision or my ability to wear contact lenses. I am aware that if I become unable to wear contact lenses while on full duty, I may be assigned to restricted duty assignments.*


*I have been informed that my use of contact lenses may be subject to verification by my employer, which may include an eye examination as necessary in the judgment of my employer's medical staff during the last month of my academy training and thereafter, unless otherwise medically indicated.*

*By my signature below, I acknowledge that I have read and accept the conditions of this Notice.*

\_\_\_\_\_  
SIGNATURE

\_\_\_\_\_  
DATE

**Figure XI-8.** Non-Compliance of LAPD Officers with Soft Contact Lenses Based on Time Since Hire (N=808)




After the first year of use, a slightly different trend appears in the general population of established lens wearers. For those who have worn SCLs for more than one year, 34%-40% reported discontinuing soft contact lens wear for more than four months, and between 12% and 24% discontinued lens wear permanently (Dumbleton et al., 2013; Pritchard et al., 1999, Richdale et al., 2007; Young et al., 2002). The mean number of years before permanent stoppage varies from 5-9 years of contact lens wear. Young et al. (2002) reported a median value of one year.

Figure XI-9 shows the most common reasons for extended lapses in established SCL wearers (Dumbleton et al., 2013) and permanent discontinuation of both established SCL and corneal RGP lens wearers (Richdale et al., 2007). Soft contact lens wearers comprised 75% of the Richdale et al. (2007) sample. The most common reason for discontinuing lens wear was ocular symptoms, such as discomfort and irritation, in both groups. Dryness and discomfort were the two most commonly reported reasons for discontinuing wear. Patients who were refitted had a short-term (i.e., 6-month) success of 77% (Young et al., 2002).

These general population SCL discontinuation rates are higher than both the 5% reported for LAPD officers during 1988-1993 and the USAF aircrew who also experienced a 5% dropout rate over 3 years (Dennis et al., 1993). One of the key differences for the lower discontinuation rate for LAPD officers, thus resulting in higher compliance, is that wearing SCLs is a condition of employment. Even successful contact lens wearers may not always wear their lenses every time they are on duty. Despite the numerous advantages of contact lenses in the cockpit (Dennis et al., 1992), only 67% of the SCL wearers and 78% of the RGP wearers wore their lenses on over 95% of the missions (Dennis et al., 1996). All the surveys indicated that permanent discontinuation of SCL wear due to medical conditions was uncommon (1/300 officers). Similar data describing the discontinuation rate for scleral lenses and HCLs after the first year of wear has not been reported.

**Figure XI-9.** The major reasons for extended lapses or permanent discontinuation of wear in primarily soft contact lens wearers. (Richdale et al., 2007; Dumbleton et al., 2013)



Taken together, studies on the discontinuation rates for SCL wearers indicate that agencies who allow a waiver for the uncorrected visual acuity requirements should monitor compliance on a continuous basis and have policies in place to manage noncompliant officers.

Scleral lenses and HCLs are an acceptable alternative to spectacles; however, it is not recommended that agencies allow a waiver for the uncorrected visual acuity requirement for these lenses because data regarding the use of these lenses in the policing environment is lacking. There is also insufficient data to determine the number of extra sick days that the officers would have to take if they could not wear their lenses, and the discontinuation rate after several years of wear is unknown. The impact of midday fogging while wearing scleral lenses on performing policing duties is also uncertain. The use of these lenses as an alternative to spectacles or SCLs, with no waiver for the uncorrected visual acuity requirements, is reasonable because the officers would still meet the visual acuity requirements if they had to remove their lenses while on duty, or they had to return to spectacle or SCL wear because they could no longer wear the scleral lenses or HCLs. As these lenses become more common in the policing environment, it may be possible to gather data to determine whether scleral lens and HCL wearers can be granted a waiver for the uncorrected visual acuity requirement.

Another important factor to consider is that the majority of scleral lens and HCL wearers have a corneal disorder or ocular surface disease (i.e., dry eye), and spectacle lenses or SCLs do not provide adequate vision (Nau, 2008; Schornack, 2017; Sarac et al., 2019; Schornack et al., 2019). Several of these disorders are progressive, and so scleral lens and HCL candidates should be questioned about corneal diseases, corneal dystrophies and corneal injuries along with the presence of any ocular surface diseases. The use of spectacles or SCL as corrective lenses does not eliminate the possibility that a candidate has a corneal or ocular surface disorder, but the probability that the candidate has some underlying ocular condition is much higher if they are wearing scleral lenses or HCLs.

Keratoconus is the most common corneal disorder encountered in the applicant pool. Keratoconus is a bilateral, progressive, noninflammatory corneal disease that results in irregular

astigmatism, central corneal thinning and corneal scarring (Kennedy et al., 1986; Wagner et al., 2007). The median age of onset is in the mid-20s and the condition usually stabilizes between the ages of 30 and 45 years (Kennedy et al., 1986; Davis et al., 2006, Hashemi et al., 2020). The prevalence of the condition varies across countries and ethnic groups. The prevalence in the United States is estimated to be 54.5/100,000, whereas prevalence reaches 3300/100,000 in some middle east countries (Ferrari & Rama, 2020). The condition affects both eyes, although the severity can vary between eyes. Both irregular astigmatism and scarring can reduce visual acuity. Historically, the cumulative rate of requiring a corneal transplant because of vision loss or contact lens intolerance was 20% over a period of 20 years after diagnosis (Kennedy et al, 1986). In an earlier study that followed previously diagnosed keratoconic patients, the cumulative rate of those patients requiring a transplant during the study's eight years was 12% (Gordon, et al., 2006). Data from more recent studies indicate that corneal RGPs, scleral lenses or corneal crosslink surgery could delay, or eliminate, the need for a corneal transplant (Godefrooij, et al, 2017; Koppen, et al., 2018; Ling, et al. 2020). In one study the number of corneal transplants over a 5-year period was reduced by 50% with the use of scleral contact lenses (Koppen, et al. 2018).

If the candidate with keratoconus (or other corneal disorders) meets the corrected and uncorrected acuity standard with spectacles, then they have the option of wearing SCLs, scleral lenses or HCLs. Spectacles could be used when they are not wearing contact lenses. However, additional information is necessary regarding the prognosis of the condition before determining whether the candidate qualifies. Since the disease is progressive, continued employment may be contingent on meeting the agency's visual acuity requirements, which may include periodic assessments of the officer's corrected and uncorrected acuities.<sup>8</sup>

It is also possible that these individuals would meet the uncorrected visual acuity standard but could only meet the corrected visual acuity standard when wearing scleral lenses or HCLs. These candidates may be acceptable provided that the wearing time of the lenses is sufficient, the condition is sufficiently stable so that they will continue to meet the uncorrected and corrected acuity requirements, and the agency is willing to accept additional sick days or reassignments for those days that the lenses cannot be worn. The number of additional sick or reassignment days is uncertain, but based on surveys of police officers who wore SCL or corneal RGPs (Hovis et al., 1998; Wells et al., 1998), the estimated median number of additional sick, or reassignment days could be 2-3 days per year.

**SUMMARY:** Soft contact lenses, scleral lenses, and hybrid contact lenses are an acceptable alternative to spectacles; however, corneal RGP contact lenses should not be permitted. Those who have worn contact lenses for less than six months should be required to meet the same corrected and uncorrected acuity as spectacle wearers. The uncorrected vision requirement may be waived for candidates who have been successful SCL wearers for at least six months. However, compliance should be monitored, and it is best ensured with a pre-placement agreement requiring officers to wear their contact lenses while on duty (see Figure XI-7). Due to a lack of data in the area, scleral and hybrid lens wearers should meet the same uncorrected far acuity standards as required for spectacle wearers. Scleral and hybrid contact lens wearers should be asked additional questions regarding corneal and ocular surface disorders. Periodic eye exams to ensure continued compliance with acuity standards may be necessary for officers with progressive eye disease.

---

<sup>8</sup>See Enforcement Guidance on Disability-Related Inquiries and Medical Examinations of Employees under the ADA (<https://www.eeoc.gov/laws/guidance/enforcement-guidance-disability-related-inquiries-and-medical-examinations-employees>)

### 3. Orthokeratology

Orthokeratology (corneal reshaping) involves the use of special RGP lenses overnight to reduce myopia by changing the shape of the cornea. It is a technique used to slow the rate of myopia progression in children and adolescents (Koffler & Sears, 2013). The procedure can enable the individual to have good vision without corrective lenses for a short period, usually 1-3 days. Two trademarked corneal reshaping procedures have FDA approval: Paragon CRT from Paragon Vision Science, and Bausch + Lomb's Vision Shaping Treatment (VST).

An individual's vision deteriorates as the cornea returns to its original shape after lens removal. In as little as one day after overnight wear, the refractive error can change by 0.25 to 0.75 diopters (Swarbrick, 2006). A 0.75 D change could reduce acuity to 20/40-20/50 range. These lenses are, therefore, not an acceptable method of correction.

Candidates who have worn corneal shaping lenses within the prior two months should be required to demonstrate that their refractive error and visual acuity have been stable for at least one month. This is not necessary for candidates whose treatment has been discontinued for two months or longer.

#### d. FAR ACUITY SUMMARY

The following recommended far acuity standards take into consideration environmental factors that may impede an officer's use of corrective lenses, the likelihood of assaults on officers occurring in low light levels, and/or the potential need for the use of lethal force. These guidelines are depicted in Table XI-12.

#### No Correction Needed

- Better Eye: **20/20**
- Weaker Eye: **20/40** for officers who use long guns or shotguns, or activities where they may need to rely on either eye separately; otherwise **20/125**.

#### Use of Spectacles, Scleral Lenses, or Hybrid Contact Lenses

- Uncorrected vision: 20/40 for *each* eye for officers who use long guns or shotguns; otherwise, 20/40 - 20/125<sup>9</sup> in the better eye and 20/125 in the weaker eye.
- Corrected vision: 20/20 in better eye; weaker eye of 20/40 for officers who use long guns or shotguns; otherwise, weaker eye of 20/125.

**Note:** To reduce the likelihood and severity of injury to the officer, spectacles that are worn on duty should consist of polycarbonate lenses and frames that meet ANSI Z87.1 specifications.

#### Use of Soft Contact Lenses

Candidates who have worn soft contact lenses (SCLs) for **less** than six months should meet the same far acuity standards as the above standards for spectacles, scleral lenses and hybrid contact lens wearers. Those who have successfully worn SCLs for longer than six months need not meet an

---

<sup>9</sup> The choice of an uncorrected standard should take into consideration the likelihood of assaults on officers, light levels, inclement weather and other environmental conditions that may affect visibility with spectacles.



uncorrected acuity standard, provided that there is a pre-placement agreement and a monitoring program in place to ensure continued use of SCLs while on duty.

The use of corneal RGP lenses is not recommended due to the risk of particle entrapment.

### Orthokeratology

Due to the potential for fluctuating vision and difficulty in ensuring compliance, orthokeratology is not an acceptable method of vision correction.

**Table XI-12.** Summary of Far Acuity Guidelines

|  | No Correction Needed                  | Spectacle Wearers, Scleral Lenses, or Hybrid Contact Lenses |                                       | SCL > 6 Months and SCL Waiver Program |                                       | SCL < 6 Months or No SCL Waiver Program        |                                       |
|--|---------------------------------------|---|---------------------------------------|---------------------------------------|---------------------------------------|--|---------------------------------------|
|  |                                       | Uncorrected   | Corrected                             | Uncorrected                           | Corrected                             | Uncorrected                                    | Corrected                             |
| <b>*Regular patrol officer duties</b>                  | 20/20 better eye<br>20/125 weaker eye | **20/40 – 20/125 better eye<br>20/125 weaker eye            | 20/20 better eye<br>20/125 weaker eye | None                                  | 20/20 better eye<br>20/125 weaker eye | **20/40-20/125 better eye<br>20/125 weaker eye | 20/20 better eye<br>20/125 weaker eye |
| <b>Duties include sighting/viewing with either eye</b> | 20/20 better eye<br>20/40 weaker eye  | 20/40 each eye  | 20/20 better eye<br>20/40 weaker eye  | None                                  | 20/20 better eye<br>20/40 weaker eye  | 20/40 each eye                                 | 20/20 better eye<br>20/40 weaker eye  |

\*Regular patrol officer duties include working in low lighting conditions, using weapons, and conditions that could result in trauma and/or spectacle dislodgement. See Patrol Officer Job Demands: Their Implications for Medical Screening.

\*\*The likelihood of assaults, inclement weather and other environmental conditions that could affect visibility (e.g., low lighting) with and without spectacles must be considered in establishing an uncorrected standard for the better eye.

### e. RECOMMENDED EVALUATION PROTOCOL

**Corrected Vision.** The most common cause of corrected vision worse than 20/20 in either eye is inadequate corrective lens prescription. However, poor corrected vision may also be indicative of serious eye disease. A vision specialist should evaluate the candidate unless the screening physician is confident that the reduction is due to childhood amblyopia.

**Uncorrected Vision.** Candidates who do not meet the uncorrected vision standard should be given an opportunity to have their vision retested by their vision specialist. Since measurement of uncorrected vision can vary with squinting, time of day, and the lighting conditions during testing, discrepancies between the results of pre-employment vision testing and the results reported by a private specialist are not uncommon. Resolving this discrepancy requires an understanding of a few basic concepts on the optics of corrective lenses:

- Lenses with a spherical shape correct either nearsightedness (myopia) or farsightedness (hyperopia). The "strength" or curvature of the lens is measured in units known as diopters (D). The diopter strength of a lens is always preceded by either a minus (-) or a plus sign (+) to denote concavity or convexity, respectively. Minus (-) lenses correct for myopia; plus (+) spherical lenses correct for hyperopia.

- Astigmatism is an optical irregularity along an axis. Cylindrical lenses aligned along the same axis can correct this error. By convention, cylindrical correction is expressed as "minus" (-) diopters, followed by the axis of the cylinder expressed in degrees.
- Eyeglass prescriptions are based on the subjective measurement of the individual's spherical and cylindrical refraction. When performed manually, this is known as the manifest refraction (MR). Although refraction can also be conducted by an automated process, it is not as accurate. Refraction is always expressed as the spherical correction, followed by the cylindrical correction. For example, -1.50 -1.00 x 90 indicates that lenses require a minus 1.5 diopter sphere combined with a 1.0 diopter cylinder aligned along an axis of 90 degrees. If there is no astigmatism, the cylinder correction is omitted. If there is only astigmatism, the spherical correction is designated as "plano" (for example, plano -4.50 x 135).

By knowing a candidate's MR, the likelihood that squinting occurred during private testing can be determined. Peter's Table (Table XI-13) can predict the most probable uncorrected distant acuity based on the manifest refraction. To use Table XI-13, find the candidate's spherical correction along the far left side of the table. If there is no astigmatism, the predicted acuity is located in the first column to the right (minus cylinders = 00). For example, if the MR is [-1.25], distant acuity is most likely 20/70.

Predicted acuity in hyperopes decreases with age. For example, an MR of [+3.00] indicates an acuity of 20/25 in a 15-year-old, but 20/200 in a 50-year-old. This age-related effect is due to the gradual loss of accommodative power of the crystalline lens in the eye. In young persons, accommodation can completely compensate for mild hyperopia.

Cylindrical correction is found along the top of the table. The axis of the cylinder can be ignored in estimating acuity. For example, [plano -2.00 x 125] = 20/70; [+1.75 -1.25 x 275] in a 28 year-old = 20/30; [-0.25 -0.75 x 50] = 20/40. Note that a small amount of astigmatism can actually improve the vision of older hyperopes. For example, a 45-year-old with an MR of [+3.00 - 2.00 x 45] is likely to have 20/80 vision, while the predicted vision of a similar hyperopic 45-year-old without astigmatism (MR of [+3.00]) is 20/200.

Astigmatism must be expressed as a "minus" cylinder when using Peter's Table. MRs written with a "plus" cylinder can be converted to minus by adding the number of cylindrical diopters to the spherical correction (axial changes can be ignored). For example, an MR of [+1.00 +1.00] is equivalent to [+2.00 -1.00]; [-1.00 +1.00] = [plano -1.00]; [-.25 +3.75] = [+3.50 -3.75].

**Table XI-13: Peter's Relationship Between Manifest Refractive Error and Uncorrected Visual Acuity**

| minus cylinders |         |        |       |       |           |        |        |        |        |         |        |         |         |        |         |         |         |
|-----------------|---------|--------|-------|-------|-----------|--------|--------|--------|--------|---------|--------|---------|---------|--------|---------|---------|---------|
| Sphere          | 00      | .25    | .50   | .75   | 1.00      | 1.25   | 1.50   | 1.75   | 2.00   | 2.25    | 2.50   | 2.75    | 3.00    | 3.25   | 3.50    | 3.75    | 4.00    |
| -2.00           | 200     | 200    | 200   | 200   | or poorer |        |        |        |        |         |        |         |         |        |         |         |         |
| -1.75           | 100     | 100    | 100   | 200   | or poorer |        |        |        |        |         |        |         |         |        |         |         |         |
| -1.50           | 80-100  | 80-100 | 100   | 100   | 100       | 200    |        |        |        |         |        |         |         |        |         |         |         |
| -1.25           | 70      | 70     | 70    | 80    | 80        | 100    | 100    | 100    | 200    |         |        |         |         |        |         |         |         |
| -1.00           | 60      | 60     | 70    | 70    | 80        | 80-100 | 100    | 100    | 100    | 100-200 | 200    |         |         |        |         |         |         |
| -0.75           | 50      | 60     | 60-70 | 70    | 70-80     | 80     | 80     | 80-100 | 200    |         |        |         |         |        |         |         |         |
| -0.50           | 30-40   | 40     | 50    | 50    | 60        | 60-70  | 70     | 80     | 80     | 100     | 100    | 100     | 100     | 200    |         |         |         |
| -0.25           | 25      | 25-30  | 30-40 | 40    | 50        | 60     | 60-70  | 70     | 80     | 80      | 100    | 100     | 100     | 100    | 100-200 | 200     |         |
| 00              | 20      | 20-25  | 25-30 | 30-40 | 40-50     | 50     | 60     | 70     | 70     | 80      | 80     | 80-100  | 100     | 100    | 100     | 200     |         |
| +0.25           | 20      | 20     | 25    | 25-30 | 30-40     | 40-50  | 50-60  | 60     | 70     | 70      | 80     | 80      | 100     | 100    | 100     | 100     | 100-200 |
| +0.50           | 20      | 20     | 20    | 25    | 30        | 40     | 50     | 50-60  | 60     | 60-70   | 70     | 80      | 80      | 100    | 100     | 100     | 100     |
| +0.75           | 20      | 20     | 20    | 25    | 25-30     | 30-40  | 40-50  | 50     | 60     | 60-70   | 70     | 70-80   | 80      | 80     | 100     | 100     | 100     |
| +1.00           | 20      | 20     | 20    | 25    | 25        | 30     | 40     | 50     | 50-60  | 60      | 70     | 70      | 80      | 80     | 80-100  | 100     | 100     |
| +1.25a          | 20      | 20     | 20    | 25    | 25-30     | 30     | 40     | 50     | 50-60  | 60      | 70     | 70      | 80      | 80     | 80-100  | 100     | 100     |
| b               | 20      | 20     | 20    | 25    | 25-30     | 30     | 40     | 50     | 50-60  | 60      | 70     | 70      | 80      | 80     | 80-100  | 100     | 100     |
| c               | 25      | 25     | 25    | 25-30 | 30        | 40     | 40-50  | 50     | 60     | 60-70   | 70     | 70-80   | 80      | 80     | 100     | 100     | 100     |
| +1.50a          | 20      | 20     | 20    | 25    | 25        | 30     | 40     | 50     | 50     | 60      | 70     | 70      | 70      | 80     | 80      | 80-100  | 100     |
| b               | 20      | 20     | 25    | 25    | 25-30     | 30     | 40     | 40-50  | 50     | 60      | 60-70  | 70      | 70-80   | 80     | 80      | 100     | 100     |
| c               | 30      | 25-30  | 25-30 | 30    | 30-40     | 40     | 50     | 50     | 60     | 60-70   | 70     | 70-80   | 80      | 80     | 100     | 100     | 100     |
| +1.75a          | 20      | 20     | 20    | 25    | 25        | 30     | 40     | 40     | 50     | 50      | 60     | 60      | 70      | 70-80  | 80      | 100     | 100     |
| b               | 25      | 25     | 25    | 25    | 25-30     | 30     | 40     | 40-50  | 50     | 60      | 60     | 70      | 70      | 80     | 80      | 100     | 100     |
| c               | 40      | 30-40  | 30    | 30-40 | 40        | 40-50  | 50     | 50-60  | 60     | 60-70   | 70     | 70-80   | 80      | 80     | 100     | 100     | 100     |
| +2.00a          | 20      | 25     | 25    | 25    | 25        | 30     | 30-40  | 40     | 50     | 50-60   | 60     | 70      | 70      | 70     | 80      | 80      | 100     |
| b               | 25      | 25     | 25    | 25    | 30        |        | 40     | 40-50  | 50     | 60      | 60     | 70      | 70      | 80     | 80      | 100     | 100     |
| c               | 50-60   | 40-50  | 40    | 40    | 40-50     | 50     | 50-60  | 60     | 60-70  | 70      | 70     | 80      | 80      | 80-100 | 100     | 100     | 100     |
| +2.25a          | 25      | 25     | 25    | 25    | 25        | 30     | 40     | 40     | 50     | 60      | 60     | 60-70   | 70      | 70     | 80      | 80      | 100     |
| b               | 25      | 25     | 25    | 25-30 | 30        | 30-40  | 40     | 40-50  | 50     | 60      | 60     | 70      | 70      | 80     | 80      | 100     | 100     |
| c               | 60-70   | 60     | 50    | 50    | 50-60     | 60     | 60-70  | 70     | 70     | 70-80   | 80     | 80      | 100     | 100    | 100     | 100     | 100-200 |
| +2.50a          | 25      | 25-30  | 25    | 25    | 25-30     | 30     | 30-40  | 40     | 50     | 50-60   | 60     | 60-70   | 70      | 70     | 80      | 80      | 100     |
| b               | 30      | 30     | 25    | 30    | 30        | 40     | 40     | 50     | 50     | 60      | 60     | 70      | 70      | 80     | 80      | 100     | 100     |
| c               | 70-80   | 70     | 60    | 60    | 60        | 60     | 60-70  | 70     | 70     | 70-80   | 80     | 80      | 80-100  | 100    | 100     | 200     | 200     |
| +2.75a          | 25      | 25     | 25    | 25    | 30        | 30     | 30-40  | 40     | 50     | 50-60   | 60     | 60-70   | 70      | 70     | 80      | 80      | 100     |
| b               | 30      | 30     | 30    | 30    | 30-40     | 40     | 40-50  | 50     | 50-60  | 60      | 60-70  | 70      | 70      | 80     | 80      | 100     | 100     |
| c               | 100     | 80     | 70-80 | 60-70 | 70        | 70     | 70     | 70-80  | 80     | 80      | 80     | 80-100  | 100     | 100    | 100     | 200     | 200     |
| +3.00a          | 25      | 25     | 25    | 25    | 30        | 30     | 30-40  | 40     | 50     | 50-60   | 60     | 70      | 70      | 70     | 80      | 80      | 100     |
| b               | 40      | 30-40  | 30    | 30-40 | 40        | 40     | 50     | 50     | 60     | 60      | 70     | 70      | 70-80   | 80     | 80      | 100     | 100     |
| c               | 200     | 100    | 80    | 70-80 | 70-80     | 80     | 80     | 80     | 80     | 80-100  | 100    | 100     | 100     | 100    | 200     | 200     | 200     |
| +3.25a          | 30      | 30     | 25    | 30    | 30        | 40     | 40     | 50     | 50     | 60      | 70     | 70      | 70      | 70-80  | 80      | 80-100  | 100     |
| b               | 40-50   | 40     | 40    | 40    | 40        | 40-50  | 50     | 50-60  | 60     | 60-70   | 70     | 70      | 80      | 80     | 80-100  | 100     | 100-200 |
| c               | 200     | 200    | 100   | 80    | 80        | 80     | 80-100 | 100    | 100    | 100     | 100    | 100-200 | 200     | 200    | 200     | 200     | 200     |
| +3.50a          | 30      | 30     | 30    | 30    | 30        | 30     | 40     | 40-50  | 50     | 60      | 60-70  | 70      | 70      | 80     | 80      | 100     | 100     |
| b               | 50      | 50     | 40-50 | 40    | 50        | 50     | 50-60  | 60     | 60     | 70      | 70     | 70-80   | 80      | 80-100 | 100     | 100     | 200     |
| c               | 200     | 200    | 200   | 100   | 100       | 100    | 100    | 100    | 100    | 100-200 | 200    | 200     | 200     | 200    | 200     | 200     | 200     |
| +3.75a          | 40      | 30-40  | 30    | 30    | 30        | 30     | 40     | 50     | 50-60  | 60      | 70     | 70      | 70-80   | 80     | 80-100  | 100     | 100     |
| b               | 60      | 50-60  | 50    | 50    | 50        | 50-60  | 60     | 60     | 70     | 70      | 70-80  | 80      | 80      | 100    | 100     | 100-200 | 200     |
| +4.00a          | 40-50   | 40     | 30    | 30    | 30        | 30-40  | 40-50  | 50     | 60     | 60-70   | 70     | 70-80   | 80      | 80-100 | 100     | 100     | 100-200 |
| b               | 70      | 60     | 50-60 | 50    | 50-60     | 60     | 60     | 70     | 70     | 70-80   | 80     | 80      | 100     | 100    | 100-200 | 200     | 200     |
| +4.25a          | 50      | 40-50  | 40    | 30    | 30        | 40     | 50     | 50-60  | 60     | 70      | 70-80  | 80      | 80      | 100    | 100     | 100-200 | 200     |
| b               | 70-80   | 70     | 60    | 60    | 50-60     | 60     | 60-70  | 70     | 70     | 70-80   | 80     | 80      | 100     | 100    | 100-200 | 200     | 200     |
| +4.50a          | 60      | 50     | 40    | 30    | 30-40     | 40-50  | 50     | 60     | 60-70  | 70      | 80     | 80      | 100     | 100    | 100-200 | 200     | 200     |
| b               | 80-100  | 70-80  | 60-70 | 60    | 70        | 70     | 70     | 80     | 80     | 80      | 100    | 100     | 100-200 | 200    | 200     | 200     | 200     |
| +4.75a          | 70      | 50-60  | 40-50 | 40    | 40        | 50     | 50-60  | 60-70  | 70     | 70-80   | 80     | 80-100  | 100     | 100    | 200     | 200     | 200     |
| b               | 100-200 | 80-100 | 70-80 | 70    | 70        | 70-80  | 80     | 80     | 80-100 | 100     | 100    | 100-200 | 200     | 200    | 200     | 200     | 200     |
| +5.00a          | 70      | 60-70  | 50    | 40    | 40-50     | 50-60  | 60     | 70     | 70-80  | 80      | 80-100 | 100     | 100     | 200    | 200     | 200     | 200     |
| b               | 200     | 100    | 80    | 70-80 | 80        | 80     | 80-100 | 100    | 100    | 100-200 | 200    | 200     | 200     | 200    | 200     | 200     | 200     |

Composite chart of refractive state to V.A. Derived from Peter's multiple tables. All figures are the denominator of the Snellen Fraction, whose numerator is 20/. Where given, a indicates age group from 5 to 15; b indicates age group from 25 to 35; c indicates age group from 45 to 55. Where not indicated, data applies to all ages. Above +3.50 sphere, acuity for c group poorer than 20/200 for all errors. From Borish, I.M., Visual Acuity, Clinical Refraction, 1970.

The following protocol for evaluating candidates whose uncorrected distance acuity is below the hiring agency's standard assumes that a recommended chart was used during the examination rather than a vision screening unit. A "line" refers to the lines on a vision chart (e.g., the 20/40 line).

**GROUP I:   UNCORRECTED ACUITY IS WORSE THAN THE AGENCY STANDARD BY ONLY ONE LINE**

These candidates should be given the opportunity to submit results from a current, private examination, using the protocol described in "General Screening Recommendations - Routine Testing," including the MR. Past records of previous eye exams should be requested, as they may reveal the candidate's true vision when not motivated to squint.

- If past records and the current private exam indicate acceptable vision, the candidate is passed.<sup>10</sup>
- If either the current private exam results *or* past records confirm unacceptable vision, the candidate should be deemed unqualified. Past records, unlike the results of a current private exam, are unlikely to be biased by squinting.<sup>11</sup>
- If the current private exam results are acceptable, but no past records are available, use the MR and Table XI-13 to assess the likelihood of squinting.

**GROUP II:   UNCORRECTED ACUITY IS WORSE THAN THE AGENCY STANDARD BY TWO LINES OR MORE**

Repeat testing by a private vision specialist is not worthwhile (unless acuity was initially screened using a vision screening unit). These candidates should be restricted from involvement in critical situations that can result in loss of spectacles. The use of soft contact lenses or refractive surgery may be an acceptable option.

***Soft Contact Lens Wearers.*** The records from the candidates' current vision specialist should be obtained (see form provided in Figure XI-10) to evaluate their experience with contact lenses and to identify possible contraindications – both absolute and relative - to successful contact lens use.

Absolute contraindications include autoimmune disorders, which are commonly complicated by the sicca syndrome (dry eyes and mouth). These include scleroderma, Sjögren's syndrome, rosacea, rheumatoid arthritis, and lupus. Diabetes is a contraindication if there are signs of ocular surface disease, a history of recurrent erosions, or corneal insensitivity (O'Donnell & Efron, 2012).

---

<sup>10</sup> This accounts for measurement error, since the standard deviation of individual differences with EDTRS or Bailey-Lovie charts on different days is approximately 3 letters and 95% of the differences fall between  $\pm 1$  line (Raasch et al, 1999; Hazel & Elliott, 2002).

<sup>11</sup> Note: Vision does not improve with age.

Figure XI-10. Sample Contact Lens Examination Form

**S A M P L E**

**CONTACT LENS DATA SHEET FOR PEACE OFFICER CANDIDATES**

**TO QUALIFY FOR THE JOB OF PEACE OFFICER, YOU MAY BE REQUIRED TO WEAR CONTACT LENSES. WE DO NOT ACCEPT USE OF HARD OR "SEMI-RIGID" LENSES DUE TO GREATER RISK OF HAVING THE LENS POP OUT OF THE EYE. PLEASE SUBMIT A CURRENT EYE EXAMINATION (WITHIN THE LAST THREE (3) MONTHS) FROM YOUR PRIVATE OPTOMETRIST OR OPHTHALMOLOGIST THAT INCLUDES ALL OF THE FOLLOWING INFORMATION:**

a. Type of contact lenses the patient currently wears: \_\_\_\_\_

b. Date patient began wearing their current contact lenses: \_\_\_\_\_

c. Average wearing time: Hours per day = \_\_\_\_\_ Days per week = \_\_\_\_\_

d. Date last pair of lenses dispensed: \_\_\_\_\_

e. Condition of current lenses: \_\_\_\_\_

f. Is there a history of any difficulties with contact lens use: \_\_\_\_\_

g. Date of last full examination of eyes: \_\_\_\_\_

h. Uncorrected distance visual acuity: OD=20/ \_\_\_\_\_ and OS=20/ \_\_\_\_\_

i. Corrected distance visual acuity with current contacts: OD=20/ \_\_\_\_\_ OS=20/ \_\_\_\_\_

j. Refractive error: OD= \_\_\_\_\_; OS= \_\_\_\_\_

k. Current spectacle prescription and visual acuities; OD= \_\_\_\_\_ 20/ \_\_\_\_\_  
OS= \_\_\_\_\_ 20/ \_\_\_\_\_

l. Please list all prescription and OTC medications: \_\_\_\_\_

m. Does the patient have any of the following conditions:

|                       |                              |
|-----------------------|------------------------------|
| _____ Corneal Ectasia | _____ Pellucid Degeneration  |
| _____ Diabetes        | _____ Rheumatoid Arthritis   |
| _____ Dry Eyes        | _____ Rosacea                |
| _____ Epilepsy        | _____ Scleroderma            |
| _____ Keratoconus     | _____ Sjögren's Syndrome     |
| _____ Lupus           | _____ Terrien's Degeneration |

n. Statement of any medical contraindication to the continued wearing of contact lenses.  
\_\_\_\_\_

o. Doctor's Printed Name: \_\_\_\_\_  
Doctor's Signature: \_\_\_\_\_  
Office Address: \_\_\_\_\_  
Phone Number: \_\_\_\_\_

Relative contraindications to contact lens use include dry eyes, use of antihistamines (which decrease tear flow), or a history of medical complications from contact lens use. Complications include corneal abrasion, corneal infection, neovascularization of the cornea (often seen in post-radial keratotomy patients who wear contacts), and giant papillary conjunctivitis (GPC). GPC is a sterile inflammatory reaction of the upper eyelid caused by friction and irritation from repetitive blinking over the upper portion of the contact lens. This condition occurs more commonly with extended wear lenses. It is often treated with steroids and temporary discontinuation of contact lens use, then refitting with a daily disposable lens.

Keratoconus, corneal ectasia, pellucid degeneration, and Terrien's degeneration are conditions that result in optical distortions due to irregularities in the corneal shape. Although scleral lenses and hybrid contact lenses are most frequently used to correct vision in these cases, SCLs may be prescribed in the early or mild cases. A positive response to any of these conditions would warrant further investigation regarding the prognosis for the stability of the refractive error, uncorrected acuities and corrected acuities.

**Corneal RGP Contact Lens Wearers.** Candidates who currently wear corneal RGP lenses should be advised to change to soft lenses, scleral lenses, or hybrid contact lenses. Those with astigmatism may require "toric" lenses. Corneal RGP lens wearers who have been refit with larger diameter SCLs can find the new lenses uncomfortable and may not adapt to the larger size and/or be unwilling to care for the lenses. Differences in material and lens size can also result in an initial change in refractive error, which will stabilize over two months. Therefore, a minimum period of two months of successful wear of the SCL, scleral lenses, or hybrid contact lenses should be required before being reevaluated.

**Scleral Lens and Hybrid Contact Lens Wearers.** Some individuals who wear these lenses have corneal or ocular surface disorder, thus it is necessary to collect additional information. The report for *all wearers* should include:

- Corrected acuities while wearing contact lenses and spectacles
- Uncorrected acuities
- Minimum, average and maximum wearing time
- Average number of days per week the lenses are worn
- Presence of midday fogging (for scleral lens wearers only)

If there is an *underlying condition* then, in addition to the above, the report must also include the diagnosis of the underlying condition and prognosis for the applicant's corrected and uncorrected acuities to continue to meet agency standards.

**Use of Contact Lenses after Refractive Surgery.** In rare cases, post-refractive individuals may require contact lenses (most likely scleral lenses or hybrid contact lenses) in order to meet the corrected acuity standard. In these cases, contact lenses are necessary to correct corneal irregularities created by the surgery. These candidates should be evaluated further to determine whether they have an increased sensitivity to glare or impaired night vision.

## 2) REFRACTIVE SURGICAL PROCEDURES

### A. General Considerations

#### 1. LASIK, LASEK, PRK, SMILE

Laser-assisted in situ keratomileusis (LASIK) and photorefractive keratectomy (PRK) are commonly-accepted procedures for correcting refractive errors. Although less common, laser

epithelial keratomileusis (LASEK) is also acceptable. The small incision lenticule extraction (SMILE) is a newer surgical option, similar to LASIK. The SMILE procedure uses a laser to make a small corneal incision in the cornea to remove some inner corneal tissue to correct the refractive error.

PRK and LASIK offer respective advantages and disadvantages. In general, LASIK and SMILE provide more rapid improvement in vision and less discomfort. PRK requires more time to recover good vision and causes more discomfort during the first weeks. The major difference between LASIK, SMILE and PRK, however, is the degree of post-surgical corneal haze. Relative to PRK, haze is less common after LASIK and SMILE (Ganesh et al., 2018). Some surgeons prefer PRK or SMILE due to concerns that the corneal flap, which is produced during LASIK, becomes dislodged due to trauma. Although there are several reports of flap dislocation due to ocular trauma up to 14 years postoperative, there are no known published reports of flap dislodgments in peace officers treated with LASIK (Patel, 2001; Iskander, 2001; Sridhar, 2001; Holt et al., 2012).

Laser refractive surgery is not without complications. The evaluation must establish that the cornea has healed, the refractive error is stable, and visual acuity meets requirements and is stable. The time required to heal and stabilize is directly related to the degree of the original refractive error (Soong & Malta, 2009; Stonecipher et al., 2006; Gil-Cazorla, 2008; Javaloy et al., 2011).

Table XI-14 reports the prevalence of post-LASIK symptoms (McDonald, 2001). The most common complaint six months after LASIK relates to driving at night: 6%-14% of patients reported that driving was significantly more difficult than before surgery. However, the study did not segregate individuals into those who obtained 20/20 uncorrected acuity from those whose uncorrected acuity ranged between 20/25-20/40. Individuals whose acuities range between 20/25-20/40 are likely (and legally) able to drive without glasses, although they do not see as well as they did prior to surgery if their corrective lenses gave them 20/20 acuity.

**Table XI-14.** Summary of (Post-LASIK) Symptoms Reported Significantly Worse at Six (6) Months<sup>a,b</sup>

|                          | <b>% Spherical Myopia<br/>n = 142</b> | <b>% Myopia with Astigmatism<br/>n = 109</b> |
|--------------------------|---------------------------------------|--|
| Night driving difficulty | 6.4                                   | 13.8   |
| Glare                    | 3.5                                   | 9.2  |
| Halos/starbursts         | 4.2                                   | 6.4  |
| Light sensitivity        | 2.8                                   | 5.5  |
| Dryness                  | 4.2                                   | 2.8  |
| Fluctuation of vision    | 2.1                                   | 1.8  |
| Blurring of vision       | 2.1                                   | 0.9  |
| Redness                  | 0.7                                   | 1.8  |
| Double vision            | 0.7                                   | 0.0  |
| Headache                 | 0.7                                   | 0.0  |

<sup>a</sup> Some subjects may represent more than one symptom.

<sup>b</sup> None of the following symptoms were reported as significantly worse: pain, burning, excessive tearing, and gritty feeling. McDonald et al. (2001)

Objective testing for glare sensitivity and contrast sensitivity can also help ensure that the essential job duties can be performed safely; however, test results need to be referenced to a nonrefractive surgery control group (Hovis & Ramaswamy, 2006).

Determination of a minimum deferral period following surgery depends on the time course of: (a), symptoms, (b) complications, and (c) regression of the surgical effect:

**a) Symptoms.** The vast majority of symptoms develop in the first week post-op (Autonomous Technologies Corp., 2000; Nidek Technologies Inc., 2000; Bausch & Lomb Surgical, Inc., 2000; Summit Technologies, Inc., 1999). Therefore, a one month deferral period should be sufficient for candidates who are symptom-free.

**b) Complications.** The vast majority of significant surgical complications are evident at one-month post-op. One exception would be ectasia or bulging of the cornea due to excessive thinning. Ectasia can develop months after surgery, but is seen primarily in patients with treatments above 10 D.

**c) Regression of surgical effect.** The time course for the regression of the surgical effect is related to the amount of correction attempted. Most individuals experience regression for only a few weeks. However, those with high degrees of myopia may progressively regress over months or years. In one FDA approval study, only 5% of patients whose pre-op manifest error were under 7 D had >1 D of regression between the 1<sup>st</sup> and 3<sup>rd</sup> months post-op. However, 12% of patients with > 7 D pre-op regressed >1 D between 3-6 months (Summit Technologies, Inc., 1999). In a study involving patients with 9-25 D pre-op refractive errors, continuous myopic regression was observed for over two years. The average regression at two years post-op was over 2.6 D (Han, 2000). In individuals with 10 D pre-op, 20% will regress by >1 D between 1-12 months post-op (Knorz, 1998).

Regression towards myopia can occur between 10-13 years post-op. As with short-term regression, the amount varies according to the type of surgery and initial refractive error. For refractive errors of less than 6 diopters of myopia before surgery, the mean change in refractive error over 10-13 years for PRK ranges from 0.46-0.64 more myopic and 0.33 for LASIK. The higher the initial refractive error, the greater the regression towards myopia (Alio et al., 2008a,b,c,d; Dirani et al., 2010). PRK eyes become 1-1.2 diopters more myopic over a 10-13 year period; LASIK eyes become 0.63-1.76 diopters more myopic. In LASIK eyes, this regression occurs mainly in the first 2-5 years (Alio et al., 2008c,d); in PRK patients, the greater rate of regression occurs within the first two years (Alio et al., 2008a,b). Except for initial refractive errors over 10 diopters, uncorrected visual acuity remains stable after two years (Alio et al., 2008b,d; Shojaei et al., 2009).

The majority of individuals who undergo laser refractive surgery to correct low to moderate refractive errors continue to meet the uncorrected visual acuity requirement of 20/40 after 10 years without requiring retreatments (Alio et al., 2008a,c; Dirani et al., 2010). However, spectacles, contact lenses, or retreatment may be needed to obtain 20/20 vision. Those with high initial refractive errors (> 6 diopters) are likely to require retreatment to attain 20/40 uncorrected and spectacles or contact lenses to achieve 20/20 corrected. Because of the risk of the refractive error regressing to unacceptable levels, annual vision testing is advised to ensure that vision remains within agency standards.

## 2. Intrastromal Corneal Rings (ICR)

In 1999, the FDA approved intrastromal corneal rings (ICR) for persons with mild myopia ( $\leq -3.00$  diopters). ICRs have the benefit of being reversible if serious side effects occur. Over time, persons treated with ICRs appear to achieve better uncorrected visual acuity than those corrected with LASIK (Suiter et al., 2000). However, the procedure has become rare except for cases with keratoconus. Intacs have been approved to manage keratoconus and



other corneal thinning disorders and diseases with good long-term success (Levinger & Pokroy, 2005; Bedi et al.; 2012).

ICR for correcting myopia requires evaluation per the general guidelines for evaluating refractive surgery, as some individuals experience problems with fluctuating acuity, glare, and double images, especially at night. Stabilization of the refractive error usually occurs by three months post-op (Schanzlin, 1999). Because ICRs are removable (removal rate during the first year is 11%), monitoring to ensure assurance that officers wear rings is necessary.

### 3. Phakic Intraocular Lenses

In 2004, phakic intraocular lens implants (PIOL) were approved to correct refractive errors too large (spherical refractive errors greater than -10.00D or + 3.50D) to be corrected by laser refractive surgery, or when laser refractive surgery is contraindicated. Three PIOLs are currently approved by the FDA: the **Artisan intraocular lens** (Ophtec USA, Boca Raton, FL), the **Verisyse** (Advanced Medical Optics, Santa Ana, CA) and the **Visian** (also known as the Implantable Collamer Lens from STAAR Surgical, Monrovia, CA). The Artisan and Verisyse are actually the same AC PIOL, distributed by two different companies. The Visian is a PC PIOL. Other PIOLs designs are in various stages of development, and once approved, will be listed at the [FDA website](#).

PIOLs can be implanted in the eye's anterior chamber (AC PIOL) or posterior chamber (PC PIOL). The surgery to implant these devices is similar to cataract surgery. PIOL surgery is primarily used to correct high degrees of myopia (Waring & Durrie, 2008). Although the PIOLs can be removed, there are no known reports of individuals doing so without precipitating complications.

PIOLs have been associated with cataract formation and potential night vision problems (glare, halo). Cataracts can be caused by surgical trauma, disruption of the aqueous flow that brings nutrients to the crystalline lens, or the crystalline lens touching the implant. Most of the cataract cases occur within three to six months, although it can take up to two years to develop. Surgically-induced cataracts that occur within the first year post-op tend to remain stable; cataracts that occur later are progressive. The incidence of surgically-induced cataracts for the Visian PC PIOL ranges from 2.5%-33%, depending on the definition of cataract and the skill of the surgeon (Lackner et al., 2003, 2004; Sanders et al., 2002, 2003, 2004). Newer-designed PIOLs coupled with more-skilled surgeons has reduced the rate of incidents to less than 3% (ITM Study Group, 2003). Most surgically-induced cataracts are mild with about half of these individuals experiencing symptoms.

Based on the PIOL studies, it is difficult to determine whether the more subtle cataracts/opacities would cause vision loss detrimental to policing. These studies report that individuals with trace focal cataracts tend to be asymptomatic; however, those with more dense cataracts report problems with glare, halos, and night vision problems. Approximately half of the individuals with cataracts after PC PIOL implantation require cataract surgery within 2-3 years. In the case of cataract surgery, the PIOL and the human crystalline lens are replaced by a single implant with generally successful results. Cataract progression is more likely for those over 40 years old, females, and those with earlier PC PIOL models. The incidence of cataract formation after one year for the approved PC PIOL model, where patient selection and surgical guidelines were followed, is less than 1% (Sanders et al., 2002; Sarikkola et al., 2005; Gimbel et al., 2018). If subtle cataracts are noted, then additional testing with respect to glare sensitivity and contrast sensitivity may be necessary along with a more detailed history about night vision problems.

The incidence of surgically-induced cataracts with AC PIOLs ranges from 2-3% (Menezo et al., 2004; Alio et al., 2002; Maloney et al., 2002). Cataracts usually appear during the first two months and remain stable. In addition to surgically induced cataracts, there is a slight regression towards myopia during the first six months after surgery (Maloney et al., 2002).

Night vision problems with both types of PIOLs occur primarily due to a small optical zone in the PIOL relative to the pupil size (Arne & Lesueur, 2000). The incidence of new night vision problems after anterior and posterior PIOLs - glare, halos and night driving difficulties – ranges from 4-12% (Arne & Lesueur, 2000; Sanders et al. 2003, 2004; ITM Study Group, 2004; Moshirfar et al., 2007b; Qasem et al., 2010). This incidence is equivalent to problems reported for refractive surgery patients.

**Effects of trauma.** There are four known reports of dislocation/dislodgement of Visian PC PIOL due to trauma (Kong et al., 2010; Schmidt et al., 2012; Espinosa-Mattar et al., 2012; Moshirfar et al., 2014). All the cases involved blunt trauma to the head or eye. One case involved dislocation due to trauma to the back of the head; the others were due to a fist striking the area around the eye. In all cases, the lens was repositioned successfully, and acuity returned to 20/20 within a week for three of the four cases. The fourth case required corneal endothelium transplantation (Kong et al., 2010; Schmidt et al., 2012; Espinosa-Mattar et al., 2012; Moshirfar et al. 2014).

Eight cases of AC PIOL dislodgement due to trauma were reported between 2002-2007 (Yoon et al., 2002; Munzo et al., 2003; Moshirfar et al., 2007a). The trauma in all cases was moderate and confined to one eye. The incidence of Artisan dislodgement is very low (near 0.1%); however, the levels of trauma were not severe – roughly equivalent to that required to displace spectacles without severely damaging an eye. Moreover, these incidents occurred in the general population, where the risk of ocular trauma was probably extremely low. Nevertheless, of the eight patients who experienced non-penetrating blunt trauma to the eye, only two patients had the AC PIOL dislodge (Moshirfar et al, 2007a).

Given the above, candidates who wear FDA-approved PIOLs and those with higher refractive errors should be deferred to ensure that: 1) the incisions have healed, 2) the refractive error is stable, and 3) cataracts are unlikely over a 12-month period. Officers with PIOLs should be required to wear protective eyewear.

#### **4. Radial Keratotomy**

Radial keratotomy (RK) has fallen out of favor due in good part to the relatively high number of complications, diurnal fluctuation in vision, (Schanzlin et al., 1986; McDonnell et al., 1996) and rupture of the globe. (Vinger et al. ,1996). Candidates contemplating RK surgery should be strongly encouraged to investigate laser refractive surgery techniques. Protective eyewear is highly recommended for officers who have had RK (Groves, 1996).

#### **B. RECOMMENDED EVALUATION PROTOCOL**

Candidates should be carefully questioned regarding problems of glare, night vision and halos. Dates of surgeries and any repeat procedures ("touch-ups" or enhancements) should be noted. Records related to the surgery and follow-up care should be obtained.

## **Laser Refractive Surgery and Intrastromal Corneal Rings (LCR)**

### **GROUP I: PRE-OP MANIFEST ERROR $\leq$ 6 D (<3 D for the ICR)**

#### **A) 1- 3 months post-operation or last enhancement:**

Acceptable if asymptomatic, refractive error has been stable, and acuity is well within acceptable limits. The refractive error is considered stable if the spherical and cylindrical components for each eye are within  $\pm 0.50$  diopter for the two assessments, separated by at least 30 days. The presence of significant symptoms or an acuity that is at the limit of acceptability would warrant deferral and reevaluation for an additional three months.

#### **B) 4-5 months post-operation or last enhancement:**

Acceptable if asymptomatic and acuity is within acceptable limits. The presence of significant symptoms would warrant deferral and re-evaluation at six months post-op.

#### **C) 6 months (or more) post-operation or last enhancement:**

Acceptable if asymptomatic and visual function is within acceptable limits. However, a history of significant symptoms at six months or longer post-op renders the candidate unsuitable, regardless of current status.

### **GROUP II: PRE-OP MANIFEST ERROR $>$ 6 D**

Acceptable if asymptomatic, visual function is within acceptable limits, and there is documentation that the manifest refraction has been stable for at least 6 months post-operative. The refractive error is considered stable if the spherical and cylindrical components for each eye are within  $\pm 0.50$  diopter for the two assessments, separated by at least 180 days. However, a history of significant symptoms at 6 months or longer post-op renders the candidate unsuitable, regardless of current status.

Due to the risk of regression, these candidates should undergo annual vision testing to ensure that visual acuity is maintained within agency standards.

## **Phakic Intraocular Lenses (PIOL)**

### **Anterior Chamber**

Acceptable if asymptomatic and visual function is within acceptable limits six months postoperative. If any lens opacities, vacuoles, or cataracts have developed, the minimum waiting period should be extended to at least six months after their first appearance.

### **Posterior Chamber**

Acceptable if asymptomatic and visual function is within acceptable limits six months postoperative. If any lens opacities, vacuoles, or cataracts have developed, the minimum waiting period should be extended to at least six months after their first appearance.

3) COLOR VISION DEFICIENCY

a. GENERAL CONSIDERATIONS

**Classification of Color Vision Deficiencies**

The human eye has three different classes of cone photoreceptors, each with a unique photopigment that preferentially absorbs different wavelengths of light (red, green, and blue). The majority of individuals with a CVD are referred to as red-green defective due to their loss of color discrimination along the red-green color axis. The major classification of CVD depends on whether: (1) one of the three distinct photopigments is different from the color-normal population in terms of which wavelengths are preferentially absorbed ("anomalous trichromats"); or (2) one of the photopigments has been replaced such that the retina contains only blue and green cones or blue and red cones ("dichromats").

CVD is further subclassified based on which pigment is involved. "Protans" have an alteration of the red receptor; "deutans" have an alteration of the green receptor. A small minority of individuals having color discrimination losses along the blue-yellow axis are referred to as "tritans." Tritans have either a nonfunctioning or partial functioning blue receptor (Table XI-15).

**Table XI-15.** Nomenclature, Classification, and Prevalence in Caucasian Males (Females) of Different Types of Human Color Vision

| Type  |             | Percentage |
|---|-------------|------------|
| <b>Trichromatic</b>                                   |             |            |
| Normal  |             | 92 (99.6)  |
| <b>Anomalous</b>                                      |             |            |
| Protan<br>(protanomalous)                             | } Red-green | 1 (0.01)   |
| Deutan<br>(deuteranomalous)                           |             | 5 (0.25)   |
| Tritan<br>(tritanomalous)                             | Blue-yellow | Trace      |
| <b>Dichromatic</b>                                    |             |            |
| Protan<br>(protanopia)                                | } Red-green | 1 (0.01)   |
| Deutan<br>(deuteranopia)                              |             | 1 (0.01)   |
| Tritan<br>(tritanopia)                                | Blue-yellow | 0.002      |
| <b>Monochromatic</b>                                  |             |            |
| S, M, or L cone<br>(incomplete or atypical achromasy) |             | 0.000001   |
| Rod<br>(typical achromasy)                            |             | 0.003      |

From Bailey, J.E. Color vision. Chapter 13 In: Clinical Procedures in Optometry. J.B. Eskridge, J.F. Amos, J.D. Bartlett (eds). Lippincott, pp. 99-120, 1991.

For the vast majority of candidates with CVD, the condition will be congenital. However, CVD can be secondary to ocular/systemic disease (such as diabetes and glaucoma) or medications (Table XI-16). Clinical characteristics that distinguish acquired CVD characteristics from those that are hereditary are presented in Table XI-17 (Bailey,1991).

**Table XI-16.** Examples of Some Commonly Prescribed Drugs Classified According to Color Deficiencies They Reportedly Induce

| Blue Defect  | Red-Green Defect  |
|--|---|
| Chloroquine<br>Indomethacin<br>Phenothiazine<br>Methimazole<br>Trimethadione | MAO-inhibitors<br>Chloramphenicol<br>Oral contraceptives<br>Ethambutol<br>Digoxin |

From Bailey, J.E. Color vision. Chapter 13 In: Clinical Procedures in Optometry. J.B. Eskridge, J.F. Amos, J.D. Bartlett (eds). Lippincott, pp. 99-120, 1991.

**Table XI-17.** Clinically Distinguishable Differences Between Acquired and Hereditary Color Vision Defects

| Hereditary   | Acquired   |
|--|--|
| Always bilateral and equal   | Usually more severe in one eye, often unilateral   |
| Almost always a red-green deficiency; much more prevalent in males | Predominantly blue-yellow defects; males and females equally susceptible; can combine with a hereditary defect |
| Other visual functions not affected                                | May affect visual acuity, visual fields, and other vision functions  |
| Stable throughout life   | Color vision varies with the status of the underlying condition; more stable if long-standing                  |
| Unambiguous color confusions on color vision tests                 | Often no clear-cut types of errors   |

From Bailey, J.E. Color vision. Chapter 13 In: Clinical Procedures in Optometry. J.B. Eskridge, J.F. Amos, J.D. Bartlett (eds). Lippincott, pp. 99-120, 1991.

### Relevance of Color Vision to Patrol Officer Duties

In the 1984 POST vision study, incumbent officers rated color identification as being "important" to "very important" (Table XI-1). Color vision was cited as being involved in an estimated 6% of critical incidents. Table XI-18 shows the frequencies for which color was used in the identification of the object.

Color identification, especially of cars and clothing, is an important component of almost all patrol officer communications. For example, when someone calls 911 and reports a suspect or vehicle, the dispatcher generally asks the caller to describe identifying colors. The subsequent radio call to a patrol car includes this information.

**Table XI-18.** Frequency of objects in the critical incidents where color was used for identification (POST, 1984).

| Object           | Percentage (N=69) |
|------------------|-------------------|
| Vehicle          | 67%               |
| Suspect clothing | 23%               |
| License plate    | 4%                |
| Container        | 3%                |
| Traffic light    | 1%                |
| Residence        | 1%                |

Steward and Cole (1989) reported that 75% of the color-defectives surveyed reported problems in selecting paints, clothes, cars, furniture and cosmetics as a result of their color vision deficiency. In a survey of 11 police officers with self-reported mild color-defects, 45% reported problems with identifying the color of cars and 18% reported problems with identifying the color of clothes (Shaw & Gledhill, 1995). In many jurisdictions, officers must be able to write legal reports and testify in court regarding their observations. A jury would likely discredit information from a CVD officer who is uncertain as to whether they saw a green car or a brown car leaving the scene of a crime, or whether a suspect had a tan or pink shirt.

Beyond color identification, color vision is also important in the recognition of signal light color. Many CVD persons report having difficulty distinguishing the color of traffic signal lights, confuse traffic lights with streetlights in the background, and have trouble seeing brake lights on cars (see Table XI-19). No CVD police officer reported problems in identifying the color of the signal light, but 9% reported difficulties in distinguishing the traffic signals from the surrounding streetlights.

**Table XI-19.** Percentage of Color Defective Individuals Reporting Difficulty with Color When Driving

| Question  | Dichromats (N=37) | Anomalous Trichromats (N=65) | Color Normals (N=102) |
|---|-------------------|------------------------------|-----------------------|
| <i>Have you ever had difficulty distinguishing the color of traffic signal lights?</i>              | 49                | 18                           | 0                     |
| <i>Do you ever confuse traffic lights with streetlights?</i>  | 33                | 31                           | 2                     |
| <i>Do you find brake lights on other cars difficult to see?</i>                                     | 22                | 8                            | 0                     |
| <i>Do you find hazard or warning lights on temporary barricades difficult to see?</i>               | 11                | 2                            | 0                     |
| <i>Do you find dashboard warning lights hard to see?</i>  | 14                | 5                            | 0                     |
| <i>Do you find some road signs such as those on freeways or school crossings difficult to read?</i> | 5                 | 11                           | 0                     |

Steward, J.M. & Cole, B.L. 1989. What do color vision defectives say about everyday tasks? *Optom. Vis. Sci.* 66(5):288-295.

The total accident rates for CVD drivers tend to be higher, although not at a level that reaches statistical significance (Verriest et al., 1980; Norman, 1980). The odds ratio for CVD drivers in

the Verriest et al. (1980) study was 1.1. Cole and Maddocks (1997) argued that there were two issues with the study. The first was the lack of statistical power. With an odds ratio of 1.1 and the assumed prevalence of CVD in the population, the sample size would have had to have been at least double in the study in order to reach statistical significance. The second issue was that Verriest et al. (1980) overestimated the prevalence of CVD in the general population. Cole and Maddocks (1997) suggest that a slightly lower rate of 7.67% is more realistic than the 8% used by Verriest et al. (1980) in calculating the odds ratio. If the lower prevalence value is used, the odds ratio of a CVD driver having an accident increases to 1.3 (Cole & Maddocks, 1997).

There is less controversy surrounding the types of accidents. Individuals who have decreased brightness sensitivity to red lights (i.e., protan defect) have twice as many accidents involving responses to red signal lights and brake lights than drivers with normal color vision, especially when conditions are wet or slippery. CVD individuals with normal brightness sensitivity (i.e., deutan defect) have twice as many accidents at intersections controlled by traffic signals, although this finding did not reach statistical significance, perhaps again due to the small sample size (Verriest et al., 1980; Cole & Maddocks, 1997).

### **Assessing Functional Abilities**

On average, dichromats have more difficulty in performing color-related tasks than anomalous trichromats, and protans generally have more difficulty driving than deutans (Verriest et al., 1980; Vingrys, 2002; Wolfe et al., 2002). Beyond these generalities, there is a wide range of functional capacity among individuals within and between all classification groups. Consequently, the primary focus of color vision testing is to first screen for color vision deficiencies before assessing their functional capacity, rather than classifying an individual's specific deficiency. The common tests include the following:

**Pseudoisochromatic Plates (PIP)** require the identification of a number or symbol consisting of colored dots embedded in a background of different colored dots. The most common PIP test is the Ishihara Color Vision Test. Different versions of the test have different numbers of test plates. If the Ishihara is used as a screening test, only the 24 plate or 38 plate editions should be administered, and the plates should be presented in random order.

Although it can quickly and accurately differentiate color "normals" from color "abnormals," the Ishihara can be compromised due to its online availability. Besides problems with test security, the Ishihara test has an overly-high sensitivity: 8% of males will fail, although not all of those who do fail will have a disqualifying degree of color vision deficiency.

A pseudoisochromatic plate test that is less subject to compromise is the Hardy Rand Rittler Color Vision Test 4<sup>th</sup> edition (HRR). The HRR consists of two parts: a screening series that distinguishes color-normals from color-defectives, and a diagnostic series that identifies the type of color vision defect and grades the severity of the defect from very mild, mild, moderate to severe. Cole et al. (2006) reported that the HRR performance is comparable to the 38-plate Ishihara test for screening red-green color vision defects if more than one error on the HRR screening series is considered a failure. Although the answers to the HRR are also readily available, the order and orientation of the test plates can be randomized, making it more difficult to memorize. To reduce the likelihood of getting the answers correct through memorization, all plates should be administered.

Computerized versions of the PIP are available. The ColorDx, for example, has sensitivity and specificity values that are nearly identical to the 38-plate edition of the Ishihara (Almustanyir, 2014; Ng et al., 2015). Not all the plates are presented once a fixed number of errors is made,

making it more difficult to memorize. Other computerized-based color vision tests should have been independently validated before being considered as a screening instrument.

**Arrangement Tests** involve arranging colored samples in a logical color sequence. The Farnsworth-Munsell D-15 (D-15) is the most widely used arrangement test. It requires an individual to place 15 colored samples (standardized paper disks mounted in caps) in a logical color sequence. The test was designed to identify CVD individuals with more severe defects who were likely to encounter problems in making color judgments encountered in ordinary occupations (Farnsworth, 1947), such as identifying color of clothes and paints. It is well standardized, readily available, inexpensive, and relatively easy to administer and score.

Table XI-20 lists the sensitivity and specificity of the D-15 for several practical color identification tasks. The general findings indicate that the sensitivity and specificity vary depending on the practical task. This variation is likely due to the number of colors to be identified, the vividness (i.e., saturation) of the sample colors and any brightness differences that may be present that provide secondary color clues. For example, in the Hovis et al. (1994) study, the sensitivity and specificity of the D-15 were greater for the small wire (i.e., low voltage applications) task than the large wire (typically 110 to 120 volt applications) task. The higher sensitivity and specificity values for the low voltage wire task were due to several factors, including smaller sized wires, the more pastel color of the small wires colors, the higher number of potential confusions in the small wires samples, and the difference in brightness relationships between the various colored small wires relative to the relationships in the large wire set. For colors used in signage, the sensitivity values generally fell between 0.70 and 0.80 (Cole & Orenstein, 2003; Cole et al., 2006).

There are two major limitations to using the D-15. First, it has low sensitivity for predicting performance on job-related duties. In 1984, POST conducted a color simulation test in which participants were shown slides and asked to name the colors of specific vehicles, suspects' clothing, traffic lights, license plates, and to determine whether vehicles' brake lights were on or off. The results indicated that persons who failed both the Ishihara test and the D-15 made significantly more errors than color normals (Table XI-21). Those who failed the Ishihara but passed the D-15 made fewer errors on all tasks than those who failed both tests; however, their error rate was almost twice that of color normals when naming the color of cars, and almost three times that of color normals when naming the color of clothing.

Individuals who receive a borderline pass on the D-15 test are more likely to have difficulty naming pastel colors (Zisman & Adams, 1985). The difference in error rates for color naming shown in Table XI-21 among color-normals, CVD individuals who pass the D-15, and CVD individuals who fail the D-15, are typical of the differences reported in other studies (Cole et al. 2003, 2006; Cole & Orenstein, 2003).

In studies performed at the City of Los Angeles, candidates who failed the Ishihara but passed the D-15 were asked to name colors of cars, clothes, buildings and paint catalog samples. Both studies found that individuals who passed the D-15 made errors in identifying colors (Goldberg, 1994, 2004). In the 2004 study, 59% (40/68) of the candidates who passed the D-15 made several errors in identifying the color of buildings and cars. Seventy-six percent (52/68) made a substantial number of errors in identifying paint chip colors. Based on these results, the ability of CVD officers who pass the D-15 to identify colors is called into question.



**Table XI-20.** Sensitivity and Specificity of the D-15 for various practical color-naming tasks.

| Study                          | Task  | Specificity<br>(Pass both D-15<br>and Practical<br>Task)                    | Sensitivity<br>(Fail both D-15<br>and Practical<br>Task)                    | Comments   |
|--------------------------------|---|---|---|--|
| Kuyk et al.<br>(1986,1987)     | Air Traffic Control.<br>Both surface colors<br>and signal lights  | 0.58  | 0.95  | Combined results for both protan<br>and deutan subjects. Mild defects<br>passed the D-15   |
| Hovis et al.<br>(1994)         | Identifying wire colors   | High voltage<br>wires: 0.72<br><br>Low voltage<br>wires: 0.93               | High voltage<br>wires: 0.73<br><br>Low voltage<br>wires: 0.85               | Low voltage wires contained more<br>colors, were smaller in size and<br>had more pastel shades   |
| Mahon &<br>Jacobs<br>(1991)    | Colors on Video<br>display (Electronic<br>Flight Information<br>System)   | Single Color<br>Presentation: 0.5<br><br>Paired Color<br>Presentation: 0.58 | Single Color<br>Presentation: 1.0<br><br>Paired Color<br>Presentation: 0.92 | Values for the single presentation<br>are based on the second series.<br>The worst normal score was the<br>cut-off score for passing the<br>practical  |
| Sui & Yap<br>(2003)            | Road markings and<br>signal lights used in<br>airports  | 0.44  | 0.88  | Limited set of surface colors (n=3)<br>and the colors and intensities of<br>the signal lights were not given   |
| Cole &<br>Orenstein<br>(2003)  | Paint, thread, fabric<br>samples<br>10 Colors<br>purple, blue, green,<br>yellow,<br>orange, brown, red,<br>black, grey and white. | Large objects: 0.68<br><br>Small objects: 0.72<br><br>Combined: 0.74        | Large objects: 0.81<br><br>Small objects: 0.74<br><br>Combined: 0.78        | Colors across materials were<br>basically the same set, but may<br>not have had the same brightness<br>relationships.<br>Visual area (degrees <sup>2</sup> ) of small<br>objects was <7 deg <sup>2</sup> and the<br>visual area of large objects was<br>>20 deg <sup>2</sup> |
| Ramaswamy<br>& Hovis<br>(2004) | Colors on Video<br>Display  | 0.82  | 0.68  | Limited to a specific railway<br>company's display   |
| Cole et al.<br>(2006)          | 10 Printed dots and<br>lines on gray<br>background<br>approximately the<br>same colors used by<br>Cole & Orenstein                | Large Sizes: 0.81<br><br>Combined: 0.92                                     | Large Sizes: 0.69<br><br>Combined: 0.70                                     | Visual area (degrees <sup>2</sup> ) of small<br>objects was <0.5 deg <sup>2</sup> and the<br>visual area of large objects was<br>>0.75 deg <sup>2</sup><br>1 error (0.8%) was allowed  |

**Table XI-21.** Color Simulation Test Results (POST, 1984)

| Color-Dependent Task | Color<br>Normals<br>(n=19)            | Fail Ishihara<br>Pass D-15<br>(n=6) | Fail<br>Both Tests<br>(n=6) |
|----------------------|---------------------------------------|-------------------------------------|-----------------------------|
|                      | <b>Number of Slides Misidentified</b> |                                     |                             |
| Color Naming:        |                                       |                                     |                             |
| Vehicles (20)*       | 4.7                                   | 8.9**                               | 11.0**                      |
| Clothing (11)        | 1.0                                   | 2.7**                               | 5.8**                       |
| License plate (5)    | 1.8                                   | 1.4                                 | 2.5                         |
| Driving-Related:     |                                       |                                     |                             |
| Brake lights (24)    | 2.7                                   | 1.8                                 | 7.4**                       |
| Traffic lights (20)  | 0.8                                   | 1.3                                 | 5.4**                       |

\*Total number of simulation slides; average number identified incorrectly is shown in table

\*\*Significantly worse than normals by t-test

The D-15 is also susceptible to improvement through practice. If shown the correct cap sequence and provided with an opportunity to practice, persons with severe color vision impairment can improve their score up to the point of passing (Ng et al., 2018a & b). Given the availability of the test both online and through purchase, the test can be compromised through practice and memorization.

In summary, unacceptable low sensitivity and the potential for test comprise seriously impact the D-15's effectiveness for determining color identification ability in peace officer candidates.

### **Hardy Rand Rittler, 4<sup>th</sup> edition (HRR) Diagnostic Plates**

Cole et al. (2007) reported that the HRR diagnostic plates were more sensitive than the D-15 in identifying individuals who named surface colors incorrectly. They and others have recommended that the predictive values of a clinical test for passing or failing the practical test are more useful indices than sensitivity and specificity because the true negatives (true positives) and false negatives (false positives) are combined into one value (Fletcher et al., 1982).

The predictive value for passing is ( $P_p$ ) is:

$$\frac{\text{Number who pass both the clinical test and practical test}}{\text{Number who pass the clinical test}}^{12}$$

The predictive value for failing is ( $P_f$ ) is:

$$\frac{\text{Number who fail both the clinical test and practical test}}{\text{Number who fail the clinical test}}$$

Table XI-22 depicts the predictive values and percent correct for various color vision tests in identifying color-defective individuals who can identify surface colors of red, orange, brown, green, yellow, blue, purple, black, white and gray based on research conducted by Cole. The studies conducted in 2007 drew from the same participant group and used the same surface colors. Cole's 2003 study used different subjects as well as colored objects that differed in color and size; moreover, the color identification task was more difficult based on the worst-normal number of errors. Percent correct represents the sum of the predictive values weighted by the probability of passing and failing the color vision test.

As indicated in Table XI-22, there is a marked similarity in percent correct values across studies. The overlap in the confidence intervals indicates that these similarities are statistically significant, consistent with Cole et al.'s 2007 finding that the predictive pass and predictive failure values for the various clinical tests were statistically identical. Notwithstanding this similarity, the HRR's higher predictive pass values indicate that it is better than the D-15 in identifying who will be able to name colors correctly. Although the HRR passed fewer individuals than the D-15, the ability of 87% of these individuals to name a set of colors is comparable to that of color normals. It is, therefore, the preferred test for screening candidates for color vision deficiencies. Although the 5<sup>th</sup> edition of the HRR test has recently been published, it has not yet been compared to the 4<sup>th</sup> edition. It is uncertain as to whether the diagnostic plates agree on the severity of the defect; therefore, it is recommended that the 4<sup>th</sup> edition be used until such data is available.

---

<sup>12</sup> Specificity is equal to number of individuals who pass both the clinical test and practical test/number who pass the practical test. Sensitivity is equal to number of individuals who fail both the clinical test and practical test/number who fail the practical test.

**Table XI-22.** Predictive values and percent correct for clinical color vision tests in identifying color-defective individuals who can identify surface colors.\*

| Study              | Test (Failure criterion)                                       | Percent who passed the clinical test (# of CVD subjects) | Predictive value for passing | Predictive value for failing | Percent correct (95% confidence interval) |
|--------------------|--|--|------------------------------|------------------------------|---|
| Cole et al. (2007) | HRR (4 <sup>th</sup> ed) (worse than mild)                     | 30 (n=99)  | 0.87                         | 0.70                         | 75.1 (65.8 to 82.6)                       |
| Cole et al. (2007) | D-15 (more than one major crossing)                            | 60 (n=99)  | 0.73                         | 0.90                         | 79.8 (70.8 to 86.5)                       |
| Cole et al. (2007) | Anomaloscope (range greater than 35)                           | 67 (n=99)  | 0.66                         | 0.97                         | 76.0 (66.7 to 83.3)                       |
| Cole et al. (2003) | D-15 (more than one major crossing)                            | 41 (n=102)   | 0.76                         | 0.73                         | 74.2 (65.9 to 82.5)                       |
| Cole et al. (2007) | Optec 900 Lantern (Farnsworth Lantern) (standard instructions) | 19 (n=99)  | 0.73                         | 0.74                         | 73.8 (65.8 to 82.6)                       |

\* red, orange, brown, green, yellow, blue, purple, black, white and gray

There appears to be limited value in administering the D-15 in addition to the HRR since the HRR has been shown to be more sensitive in identifying milder forms of color vision defects. Almoustanyir (2014) found that only 17% of CVDs who passed the D-15 (n=18) were classified as mild or very mild by the HRR, and no one who failed the D-15 (n=29) was classified as mild or very mild. These data indicate that situations in which a candidate fails the D-15 but is classified as mild by HRR should be treated with suspicion.

### Additional Tests

**The Farnsworth Munsell 100 Hue test (FM-100)** is one of the few clinical tests that can measure fine color discrimination of both color-normal and color-defective individuals. It is commonly used to assess the color discrimination ability of quality control personnel and in acquired color vision defects. Using a total error score greater than 90 as a failure, the FM-100 performs better than the D-15 in predicting performance on the color wire identification task (Hovis et al., 1994). However, the recommended cut-off score rises for individuals older than 25 years old, particularly for those in their late 40's and 50's, thereby making it a poorer predictor than the D-15 for those age groups.

The sensitivity of FM-100 for identifying high voltage wire colors is 1.0, with a specificity of 0.79. For low voltage wire color identification, the sensitivity remains at 1.0 with a specificity of 0.91 (Hovis et al., 1994). Although it may also be useful in assessing a candidate's color vision with borderline results on other color vision tests; the FM-100 requires 30-60 minutes to administer and score.

**Lantern Tests** such as the Farnsworth Lantern test (replaced by the Optec 900), require the identification of small colored lights. These tests are commonly used to certify pilots and ship captains (Hackman & Holtzman, 1992). Some have advocated their use in determining whether CVD individuals qualify for commercial driving licenses (Cole, 1993). Practical problems with the test include the limited availability of testing equipment and a testing protocol that requires modification due to problems with the accompanying instructions (Cole et al., 2008). The D-15 is slightly easier than the Farnsworth/Optec Lantern; approximately 60% of the CVD individuals who pass the D15 can pass the lantern. Of those that fail the D-15, 90% fail the lantern test (using the traditional testing protocol) (Cole & Vingrys, 1983).

**The Anomaloscope** uses color matching to identify and classify red-green color vision. Candidates adjust the relative amounts of red and green lights until they match a yellow reference standard. The severity of the defect is determined by the range of red-green mixtures that appear identical to the reference yellow light. A range  $\leq 35$  units would be considered acceptable (Cole et al., 2007).

The Anomaloscope and Optec 900 Lantern test may be of value in more challenging cases. These tests are more likely to be found at Schools and Colleges of Optometry or Departments of Ophthalmology. The Optec scoring criterion should follow the recommendation by Cole et al. (2008) of one practice run, followed by two runs of the nine pairs. The passing criterion should be no more than one total error (each misnamed light is a mistake) on the last two runs.

As discussed above, the HRR diagnostic is preferable to the Farnsworth D-15, Farnsworth-Munsell 100 Hue, Farnsworth Lantern (or Optec 900), and Anomaloscope tests in assessing the ability to determine a candidate's ability to name surface colors (see Table XI-22). Nevertheless, other test results may be useful in borderline cases or in cases where there is a suspicion that the HRR test was compromised. The caveat is that the Farnsworth D-15 could also be compromised; thus, it may be necessary to administer additional tests to confirm the Farnsworth D-15 results or select another test.

**SUMMARY:** Color vision is critical in performing a variety of job functions (e.g., identification of cars, clothing, identifying traffic lights, distinguishing traffic lights from streetlights, etc.). However, candidates with very mild or mild color vision deficiencies possess sufficient color identification and color discrimination ability. The HRR is the most effective test in distinguishing those with mild deficiencies from candidates who have more severe forms of CVD.

### **Color Corrective Lenses**

Colored contact lenses or spectacles (i.e., X-Chrom, X-Chrome, Colormax, Enchroma, Chromagen, Vino) introduce both brightness and color differences between the figure and the background colors. The effect is similar to looking at the test figures through a red filter. This violates the basic illumination requirements for any pseudoisochromatic plate test, including the Ishihara, D-15, and HRR. Performance on other color vision tests may be worse and discrimination of colors not previously confused may be poorer (Matsumoto et al., 1983; Kassir et al., 1984; Hovis, 1997; Swarbrick et al., 2001; Moreland et al., 2010).

Functional ability can be adversely affected by color correction lenses as well. If a single lens is worn, depth perception may be impaired (Matsumoto et al., 1983). Any "corrective lens" has approximately the same luminous transmittance as sunglasses (Hovis, 1997; Moreland et al., 2010); therefore, these lenses cannot be worn at night. Moreover, some lenses may make traffic light signals and LED information signs difficult to detect when worn in front of both eyes during the daytime. For these reasons, "color corrective" spectacles or contact lenses are not acceptable devices to aid in color discrimination.

### **b. RECOMMENDED EVALUATION PROTOCOL**

**HISTORY** - Table XI-23 lists the common problems encountered by color-defectives in their daily lives and the percentages responding "yes" to each question. The first 5-7 questions can serve as the basis for determining whether there is a disqualifying color vision problem based on the relatively high percentages of CVDs who encounter these problems. However, the reverse is not true: a failure to acknowledge problems does not negate the findings of objective testing, especially in high-stakes situations such as employee screening. Steward and Cole (1989) found that 5% of dichromats and 25% of anomalous trichromats were not aware of their CVD. In certain cases, consideration may be given to whether the CVD is acquired and potentially reversible (see

Tables XI-16 & XI-17). This is especially important if the CVD candidate is female, taking medication, or if the deficiency follows a tritan pattern.

**Table XI-23:** Percentage of Individuals Reporting Difficulty with Everyday Tasks That Involve Color

| Question  | Dichromats (N=37) | Anomalous Trichromats (N=65) | Color Normals (N=102) |
|---|-------------------|------------------------------|-----------------------|
| <i>Have you ever had any difficulty in selecting the colors of clothes, accessories, cars, paints, carpets, furniture, wallpaper, or cosmetics?</i> | 86**              | 66*                          | 0                     |
| <i>With craftwork and hobbies, do you have any trouble distinguishing the colors of wires, threads, materials, wools, paints, or other things?</i>  | 68***             | 23***                        | 0                     |
| <i>Do you find plant or flower identification difficult because of color?</i>   | 57***             | 18***                        | 0                     |
| <i>Have you ever had difficulty distinguishing the color of traffic signal lights?</i>  | 49**              | 18*                          | 0                     |
| <i>Do you have any difficulty determining when fruits and vegetables are ripe by their color?</i>   | 41*               | 22*                          | 0                     |
| <i>Can you determine if meat is cooked by its color?</i>  | 35*               | 17*                          | 0                     |
| <i>Do you ever confuse traffic lights with streetlights?</i>  | 33                | 31                           | 2                     |
| <i>Do you have any difficulties because of color as either a spectator or participant in sporting activities?</i>                                   | 32                | 18                           | 0                     |
| <i>Do you find it difficult to adjust the color balance on a color TV satisfactorily?</i>   | 27                | 18                           | 2                     |
| <i>Have you ever had difficulty in recognizing skin conditions such as sunburn and rashes?</i>  | 27                | 11                           | 0                     |
| <i>Do you find brake lights on other cars difficult to see?</i>   | 22                | 8                            | 0                     |
| <i>Do you find dashboard warning lights hard to see?</i>  | 14                | 5                            | 0                     |
| <i>Do you find hazard or warning lights on temporary barricades difficult to see?</i>   | 11                | 2                            | 0                     |
| <i>Do you find some road signs such as those on freeways or school crossings difficult to read?</i>   | 5                 | 11                           | 0                     |
| <i>Have you ever taken the wrong tablet or medicine because of difficulties with its color?</i>   | 0                 | 3                            | 0                     |

Significant difference between dichromats and anomalous trichromats at \*p < 0.05, \*\*p < 0.01, or at \*\*\*p < 0.002 using Yates  $\chi^2$ . From Steward & Cole, 1989. What do color vision defectives say about everyday tasks? *Optom. Vis. Sci.* 66(5):288-295.

### Set-up of Testing Area

The only light source in the test room should be the illumination provided by a True Daylight Illuminator (with slant easel) utilizing a single Verilux F15T8VLX 15w tube [or tube of equivalent correlated color temperature (between 5000 K and 7000 K and color rendering index (at least 90))]. Any other sources of extraneous light should be eliminated.

## Pre-Testing Procedures

Prior to a candidate entering the testing room, carefully inspect the subject's eyes to ensure that colored contact lenses or spectacle lenses (such as a red X-Chrom or Enchroma) are not worn.

Seat the patient so that their eyes are about 30 inches from the test book when it is placed on the easel of the Illuminator.

## HRR Pre-Testing Procedures

Demonstration Plates (1-4):

Show the first plate in the sequence and say:

*The test itself is made up of just these three symbols, an "X", an "O", or a "Δ", with two symbols, one symbol, or no symbols on a page. Some of them will be harder for you to see as they may be less strong in color.*

Open the book to the first demonstration plate. *"Here are examples of a circle and X"*

Show the second plate and state: *"Here are examples of a triangle and X"*.

Show the demonstration third plate and state:

*"Here is a circle. There can be one or two figures, or (as you turn to 4<sup>th</sup> plate) no figure. You need to point to the location of each figure that you see with the paintbrush. If you are not sure, trace the figure with the paintbrush."*

*It is important for you to tell me immediately how many symbols you see and where they are located on each page. You cannot change your answer. If you do not give me an answer within 3 seconds, I will have to turn the page.*

*"Do you understand the instructions?" Do you have any questions?"*

Test Plates:

**Due to the limited number of screening plates, it is possible to memorize the correct answer for each. For this reason, all the plates should be presented, and the order should be randomized. Each plate should be rotated so that the figures are in a different location from the one on the score sheet. Careful recording of results is critical.**

*OK, we are now ready to start the test."*

(Turn to the first test plate and proceed.)

*"What colored symbols do you see here?"*

*"Locate them with the brush."*

If the candidate responds within 3 seconds, record the response (X, O, Δ) in the box provided on the customized scoring sheet, recording the exact symbols seen in the location indicated by the candidate. If no response is given within approximately 3 seconds, mark the missed figures as errors and turn to the next test plate.

Score the test as previously described. A very mild or mild color vision deficiency is acceptable. Moderate to severe color vision deficiencies are disqualifying.

#### 4) OTHER VISUAL FUNCTIONS

##### **Visual Field Deficiency**

###### a. GENERAL CONSIDERATIONS

Partial loss of visual field in one or both eyes affects about 3% of the population between the ages of 16 to 60 (Johnson & Keltner, 1983). This increases to about 6% between the ages of 61 to 65, and 13% in persons over the age of 65. A large number of eye conditions can cause loss of visual field, the most common being glaucoma.

The 1984 POST vision survey indicated that peripheral vision is one of the most important visual abilities for safe patrol officer performance (Table XI-1). Examples of critical situations in which peripheral vision is important include:

- a suspect approaching the officer from the far right or left side;
- a hostile crowd surrounding an officer;
- an officer attempting to look out of the side of a patrol car to spot a suspect while still controlling the vehicle;
- driving under emergency conditions.

Several studies have examined the performance of persons with visual field defects in situations similar to those cited above. Johnson et al. (1992) tested the impact of spectacles that restrict peripheral vision on the ability of a correctional officer to detect suspicious behavior by inmates gathered in a day room. Restricting the horizontal binocular field to 120 degrees in each eye had no impact, but further restriction to 60 degrees significantly impaired performance, especially in detecting more subtle behavior, such as hand/arm movements and inmates reaching for the door.

In a scenario in which a parole officer was interviewing a parolee in his residence, the parole officer's ability to detect another person walking into the room or a person peaking around the corner dropped from nearly 100% for full binocular visual fields to 40% when the field was restricted to 120°. Correct responses dropped to less than 10% when the visual field was restricted even further (Johnson & Brintz, 1996).

Good et al. (2005) reported that individuals with monocular vision missed 11.4% of lighted targets overall compared to their full binocular visual field performance, even though eye, head and body movements were allowed. The difference between monocular and binocular performance was greater for the shorter duration flashes, with 16.7% more misses monocularly for the 0.1 sec flashes and 8.6% more misses monocularly for the 0.8 sec flashes.

**Driving and Visual Field Defects.** Johnson and Keltner (1983) found only slightly higher accident rates among drivers with unilateral field defects or monocularly. The drivers' visual defects were rated as severe in only 13% of the drivers with unilateral defects. However, the results of most studies that focus on monocular drivers or those with gross reductions of the visual field on one side paint a different picture. Kite and King (1961) observed a seven-fold

increase in intersection crashes and pedestrian injuries. Keeney (1968) found that monocular vision was four times more common in those cited for multiple driving violations. A pathology study concluded that long-standing lesions, which likely produced visual field defects, contributed to the death of two drivers and five pedestrians in Maryland (Freytag & Sachs, 1969).

Johnson and Keltner (1983) found that accident and conviction rates of drivers with visual field loss in both eyes were more than twice as high as those with normal visual fields. This finding is consistent with that of Hedin and Lovsund (1987), who evaluated individuals in a driving simulator. Eighty-five percent of the 27 patients with a variety of field defects had significantly increased reaction times to stimuli presented in visual areas of relevance to traffic safety. Despite the participants' ability to move their heads during testing, only 4 (15%) could compensate for their field defects.

Results from general driving field trials of those with bilateral visual losses are consistent with the previous studies. Wood et al. (2009) and Kasneci et al. (2014) reported that 25%-40% of drivers with bilateral quadrantanopsia or hemianopsias were judged as unsafe. They had problems maintaining lane position and in gap judgment ability. In the Kasneci et al. study (conducted in the United States), the majority of those who failed had left hemianopsias (visual field loss on the left side of the fixation point in each eye).

Similar peripheral vision standards were upheld in a 1988 case heard by the California Fair Employment and Housing Commission involving a monocular police officer candidate (DFEH v. City of Merced PD, FEP85-86, 88-20). In finding for the city, the Commission agreed that "peripheral vision is among the most important visual abilities that a police officer needs to safely fulfill his or her duties," and that safety concerns were not mitigated by that candidate's seven years of prior experience as a patrol officer.

**SUMMARY:** Monocular vision or significant bilateral field defects threaten personal and public health and safety. Peripheral field losses impair driving, the ability to maintain situational awareness and to detect potential hazards when monitoring a scene or when interviewing a person in the field. Significant field defects include monocular vision and cases in which the total horizontal field is restricted to < 120 degrees in either eye, the total vertical field is less than 100 degrees, or when large scotomas are present.

#### b. RECOMMENDED EVALUATION PROTOCOL

Routine screening techniques for visual field defects have poor sensitivity and specificity. Confrontation visual field testing has a sensitivity of only 37% (Johnson & Baloh, 1991), which is not surprising, given that the technique is unsuitable for detecting small patchy defects and accurate scotomas. Sensitivity for large defects such as homonymous hemianopia and altitudinal defects is 81%.

Formal perimetry testing should be conducted on those with either a personal or family history of glaucoma, any personal history of visual problems related retinal/optic nerve diseases or disorders, or decreased visual acuity (worse than 20/40) in either eye, which cannot be corrected with lenses and was not previously diagnosed as amblyopia. History of severe head trauma or stroke also indicates formal testing. Candidates with monocular vision, <120 degrees of total horizontal field in either eye, <100 degrees of vertical field, or significant scotoma are not acceptable.

The required visual field tests will depend on the specific case; however, a Humphrey's Full Field 120 screening test (or equivalent) for each eye should be included in every evaluation.



## **Binocular Fusion Deficiency**

### **a. GENERAL CONSIDERATIONS**

Abnormal binocular vision can occur when there is strabismus, and the eyes are directed at different points in the environment. The eye may be intermittently or constantly turned inward (esotropia), outward (exotropia), or vertically deviated (hypertropia). Strabismus is observed in about 4-7% of children. In childhood-onset strabismus, the resulting diplopia and visual confusion become the stimuli for suppression of the deviated eye; if not treated at a young age, this may result in permanent loss of vision in the deviated eye (amblyopia). In the case of adult-onset strabismus, suppression rarely develops, and therefore double vision is a continuing problem.

Diplopia (i.e., double vision) occurs when there is a misalignment of the eyes. It can be constant or intermittent. It is more likely to occur in reduced visual environments (i.e., nighttime) or when the individual is tired. If the diplopia is intermittent, the individual can simply close one eye; however, this reduces the field of vision and depth perception, and the duration of the diplopia is difficult to predict.

Diplopia is incompatible with many, if not most, law enforcement activities. For example, 82% (14/17) of patients with strabismus who experienced diplopia reported difficulty in driving: specific comments were “driving is terrible,” “extra caution when driving” and “afraid of hitting something on the side of the road” (Hatt et al., 2007).

Strabismus can also result in a reduction in the binocular visual field. This occurs in individuals who have large-angle esotropia (>5 degrees or 10 prism diopters) because the monocular temporal field crescent in the turned eye now overlaps with the nondeviating eye’s visual field (Kushner, 1994; Quah & Kaye, 2004). The reduction in the extent of the visual field is approximately 13% on average as compared to individuals without strabismus (Quah & Kaye, 2004), but the difference could be larger for individuals with larger amounts of esotropia (Kushner, 1994).

**SUMMARY:** The risk of diplopia and the restriction of the binocular visual fields in the cases of large angle esotropia is a contraindication to job performance.

### **b. RECOMMENDED EVALUATION PROTOCOL**

Candidates with a difference of at least two lines in their corrected monocular acuities or a history of strabismus or lazy eye should be referred to a vision specialist for further assessment to determine whether double vision is likely under both day and night conditions. Several tests are available for this purpose, such as the Worth 4 dot, red lens test and Bagaloni lenses. These tests should be administered in both light and dark, and in the six cardinal positions of gaze (straight right, straight left, upper right/left and lower right/left). A significant esotropia (>10 prism diopters) is a disqualifying reduction in binocular visual field. One automated visual field test that can assess the binocular visual field is the Esterman Binocular protocol. The total extent in the horizontal meridian should be at least 140 degrees.

### **Ancillary Tests**

**Contrast Sensitivity.** Peace officer duties depend on both visual acuity and contrast sensitivity - the ability to rapidly and accurately detect and identify objects that range in size and contrast with the background. Examples of particularly challenging visual tasks include detecting a dark-

colored weapon lying on an asphalt-covered road or identifying a dark-colored object in a suspect's hand, held closely to a dark-colored jacket.

Standard testing of visual acuity involves the identification of a black letter on a white background. The size of the letter is varied, while the contrast between the letter and the background is held constant. Given that the real world consists of more complex targets that vary in both size and contrast, both parameters are critical to the visibility of an object.

Contrast sensitivity can be measured in two general ways. One method is to maintain a constant letter/grating (i.e., alternating black and white bars) size and vary the contrast until the pattern is no longer visible. This is repeated for different sized patterns (i.e., spatial frequencies). Two clinical charts are available that have a fixed letter size and reduce the contrast until the letter is no longer visible. The Pelli-Robson chart has large-sized letters to measure low spatial frequency contrast sensitivity and the Small Letter Acuity Charts measure contrast sensitivity to high spatial frequencies (Rabin, 1994).

**Low Contrast Letter Acuity.** Charts that measure visual acuity using low contrast letters include low contrast versions of the Bailey-Lovie chart, ETDRS, and Regan Charts. In conjunction with the high contrast visual acuity, the low contrast acuity measure is often used as a surrogate for measuring contrast sensitivity.

**Glare Testing.** Glare sensitivity is normally measured by repeating some type of visual resolution test with glare light sources slightly off the direct line of sight. Commercial glare testing equipment is available from [Vectorvision](#) or the Mesotest IIb from [Oculus](#). These tests may be useful in assessing glare sensitivity in post-refractive surgery individuals and in those who have cataracts but relatively good acuity.

**Low Light Level Acuity.** Low light level acuity is measured by repeating a visual resolution test while wearing filters that reduce the amount of light entering the eyes. The filters are selected for a specific condition or are standard with a commercial test (Miller et al., 2005; Hovis & Ramaswamy, 2006). It is notable that officers interviewed in the 1984 POST vision study rated dark adaptation as the most important visual skill used on the job (see Table XI-1). However, based on officer interviews in Canada, this skill may involve seeing and detecting objects in low light levels rather than dark adaptation per se.

Contrast sensitivity has been found to correlate with performance in detecting aircraft, flight performance, simulated aircraft crash sites, object recognition and military vehicle recognition (Ginsberg et al., 1982; Stager & Hameluck, 1986; Shinar & Gilead, 1987); however, the results were not always replicable within or across studies (Kruk & Regan, 1983; O'Neal & Miller, 1987; Temme et al., 1991). The lack of repeatability could be due to the use of different equipment and procedures to measure acuity and contrast sensitivity. Studies that have used a finer scale for both measurements were less likely to find correlations between performance and contrast sensitivity (Kruk & Regan, 1983; O'Neal & Miller, 1987; Temme et al., 1991).

Routine testing on candidates who are visually normal and have a visual acuity of 20/20 or better is not necessary. However, testing contrast sensitivity, low contrast acuity, low light level acuity, or glare testing can be useful for candidates whose corrected or uncorrected acuity is borderline. This could be due to corneal irregularities, mild cataracts, out-of-date spectacle or contact lens prescriptions or contact lenses with moderate-to-heavy deposits on the surface. Their contrast sensitivity or glare sensitivity could be impaired.

Testing contrast sensitivity on all refractive surgery candidates is inefficient. Less than 10% of officers and candidates who undergo refractive surgery and have 20/20 acuity would have

unacceptable low contrast acuity and glare sensitivity. Only 15% have unacceptable low contrast acuity in low light levels (Hovis & Ramaswamy, 2006). Only a minority of the post-refractive surgery candidates have enhanced glare sensitivity or reduced vision in low light levels (Jacobs et al., 1998; Miller et al., 2005).

Several risk factors increase the probability of impaired low light level acuity or glare sensitivity. These include borderline acuity results on typical clinic acuity charts, only one eye at 20/20 or better, vague symptoms of glare sensitivity, original refractive error greater than 6 diopters, corneal irregularities, radial keratotomy, post-surgical complications, or a persistent mild haze in the central cornea.

A candidate's results must be interpreted relative to an age and acuity matched normative database. In one study (Hovis & Ramaswamy, 2006), low contrast acuity was measured using the Bailey-Lovie low contrast letter chart while wearing filters that transmitted approximately 1% of the light. This filter equates the light levels of an urban environment at night. Individuals with low contrast acuity worse than 20/80 (0.62 logMAR) were unable to read license plates from a safe distance. The 20/80 cut-off was three standard deviations from the mean acuity value of police officers and recruits who did not undergo refractive surgery.

## SUMMARY OF VISION GUIDELINES

The vision guidelines summarized below assume that: (1) officers may work in areas where environmental factors impede the use of corrective lenses; (2) assaults on officers may occur in low light levels; and (3) the use of lethal force may be required. It is important to consider the relevance of these duties and conditions when developing agency-specific vision standards.

### 1) FAR ACUITY<sup>13</sup>

The following far acuity guidelines are a repeat of the Far Acuity Summary pp. XI-34–35. Table XI-12 provides a summary of the far acuity guidelines.

#### No Correction Needed

- Better Eye: 20/20
- Worse Eye: 20/40 for officers whose responsibilities include the use of long guns or shotguns or activities where they may need to rely on either eye separately; otherwise, 20/125.

#### Use of Spectacles, Scleral Lenses, or Hybrid Contact Lenses

- uncorrected vision of 20/40 for *each* eye for officers who are required to use long guns or shotguns; otherwise 20/40 - 20/125<sup>14</sup> in the better eye and 20/125 in the worse eye
- corrected vision of 20/20 in better eye; 20/40 in the weaker eye for officers who are required to use long guns or shotguns; otherwise 20/125 in the weaker eye

**Note:** To reduce the likelihood and severity of injury to the officer, all spectacles worn by officers on duty should consist of polycarbonate lenses and frames that meet ANSI Z87.1 specifications.

#### Use of Soft Contact Lenses

Candidates who have worn soft contact lenses (SCLs) for less than six months should meet the same far acuity standards established for spectacle, scleral and/or hybrid lens wearers. However, candidates who have successfully worn SCLs for longer than six months need not be required to meet an uncorrected acuity standard, provided that there is a program in place to ensure continued the use of SCLs while on duty (i.e., pre-placement agreements with monitoring).

#### Orthokeratology

Due to the potential for fluctuating vision during and across days and the difficulty in establishing protocols for monitoring compliance, orthokeratology is not an acceptable method of vision correction for peace officers.

---

<sup>13</sup> The use of standardized charts and methods when measuring visual acuity is critical. Non-standardized testing results in erroneous measurements and increased measurement variability.

<sup>14</sup> The choice of an uncorrected standard should take into consideration the likelihood of assaults on officers, light levels, inclement weather and other environmental conditions that may affect visibility with spectacles.

## **2) REFRACTIVE SURGICAL PROCEDURES**

### **Laser Refractive Surgery (LASIK, LASEK, PRK, SMILE) and Intrastromal Corneal Rings (ICR)**

Pre-Op Manifest Error  $\leq 6$  D ( $<3$  D for the ICR)

A) 1- 3 months post-operation or last enhancement:

Acceptable if asymptomatic, refractive error has been stable for at least one month, and acuity is well within acceptable limits. The presence of significant symptoms or acuity that is at the limit of acceptability would warrant deferral and reevaluation after three months.

B) 4-5 months post-operation or last enhancement:

Acceptable if asymptomatic and acuity is within acceptable limits. The presence of significant symptoms would warrant deferral and re-evaluation at 6 months post-op.

C) 6 months (or more) post-operation or last enhancement:

Acceptable if asymptomatic and visual function is within acceptable limits. However, a history of significant symptoms at or beyond 6 months post-op would warrant disqualification, regardless of current status.

Pre-Op Manifest Error  $> 6$  D

Acceptable if asymptomatic, visual function is within acceptable limits, and there is documentation that the manifest refraction has not changed by more than 0.5 D over the last six post-operative months. However, a history of significant symptoms, at or beyond six months post-op, would warrant disqualification, regardless of current status.

Due to the risk of regression, these candidates should undergo annual vision testing to ensure that visual acuity is maintained within agency standards.

### **Phakic Intraocular Lenses (PIOL)**

Anterior Chamber

Acceptable if asymptomatic and visual function is within acceptable limits six months post-operative. If any lens opacities/vacuoles/cataracts have developed, the minimum waiting period should be extended to at least six months after the first appearance of the opacities/vacuoles/cataracts.

Posterior Chamber

Acceptable if asymptomatic and visual function is within acceptable limits six months post-operative. If any lens opacities/vacuoles/cataracts have developed, the minimum waiting period should be extended to at least six months after the first appearance of the opacities/vacuoles/cataracts.

### **Radial Keratotomy**

Radial keratotomy (RK) is less common, due in good part to the relatively high number of complications, diurnal fluctuation in vision, and rupture of the globe. Candidates contemplating RK

surgery should be strongly encouraged to investigate newer laser refractive surgery techniques. Protective eyewear should be worn by officers who have had RK.

### **3) COLOR VISION DEFICIENCY**

Those with a very mild or mild color vision deficiency as identified on the HRR have sufficient color identification and color discrimination abilities. Greater degrees of color vision deficiency are unacceptable.

Use of colored contact lens or spectacles (i.e., X-Chrom, X-Chrome, Colormax, Enchroma, Chromagen, Vino) should not be permitted during testing.

### **4) OTHER VISUAL FUNCTIONS**

#### **VISUAL FIELDS**

Candidates with monocular vision, <120 degrees of total horizontal field in either eye, <100 degrees of vertical field, or significant scotoma should be restricted from field duty. However, a monocular scotoma that overlaps onto a normal visual field in the other eye may be acceptable.

A significant scotoma is defined as an area extending more than 5 degrees in any direction where sensitivity is less than 10 dB.

#### **Binocular Fusion Deficiency**

Candidates with a difference of at least two lines in their corrected monocular acuities or a history of strabismus or lazy eye should be referred to a vision specialist for further assessment to determine whether double vision is likely under both day and night conditions. A significant esotropia (>10 prism diopter) constitutes a disqualifying reduction in binocular visual field. The total extent in the horizontal meridian should be at least 140 degrees.

#### **Contrast Sensitivity, Low Light Acuity and Glare**

Testing candidates who are visually normal and have a visual acuity of 20/20 or better is unlikely to be helpful. Testing contrast sensitivity, low contrast acuity, low light level acuity, or glare testing can be useful for candidates who are borderline with respect to meeting the corrected or uncorrected acuity requirement, as well as those with a history of corneal disorders, cataracts, retinal disease or disorders, and optic pathway diseases and disorders, even if their acuity meets the requirements.

For positions in which a large percentage of work is carried out in low light conditions, low contrast visual acuity testing with dim lighting (simulated by using tinted lenses that transmit 1% of the light) may be used for the above candidates and candidates who have had refractive surgery. Although results must be interpreted relative to an age and acuity matched normative database, individuals with a low contrast acuity in dim lighting worse than 20/80 (0.62 logMAR) may be unable to read license plates from a safe distance.

## REFERENCES

- Abdalla, Y. F., Elsahn, A. F., Hammersmith, K. M., & Cohen, E. J. (2010). SynergEyes Lenses for Keratoconus. *Cornea*, 29(1), 5-8.
- Abou Samra, W. A., El-Emam, D. S., & Kasem, M. A. (2018). Clinical Performance of a Spherical Hybrid Lens Design in High Regular Astigmatism. *Eye & Contact Lens: Science & Clinical Practice*, 44, S66-S70.
- Alió, J.L., Muftuoglu, O., Ortiz, D., Artola, A., Pérez-Santonja, J.J., de Luna, G.C., Abu Mustafa, S.K. & Garcia, M.J. (2008a). Ten year follow-up of photorefractive keratectomy for myopia of less than 6 diopters. *Am J Ophthal*, 145, 29-36.
- Alió, J.L., Muftuoglu, O., Ortiz, D., Artola, A., Pérez-Santonja, J.J., de Luna, G.C., Abu Mustafa, S.K. & Garcia, M.J. (2008b). Ten-year follow-up of photorefractive keratectomy for myopia of more than 6 diopters. *Am J Ophthal*, 145, 37-45.
- Alió, J.L., Muftuoglu, O., Ortiz, D., Pérez-Santonja, J.J., Artola, A., Ayala, M.J., Garcia, M.J. & de Luna, G.C. (2008c). Ten-year follow-up of laser in situ keratomileusis for myopia of up to 10 diopters. *Am J Ophthal*, 145, 46-54.
- Alió, J.L., Muftuoglu, O., Ortiz, D., Pérez-Santonja, J.J., Artola, A., Ayala, M.J., Garcia, M.J. & de Luna, G.C. (2008d). Ten-year follow-up of laser in situ keratomileusis for high myopia. *Am J Ophthal*, 145, 55-64.
- Alió, J.L., Mulet, M.E. & Shalaby, A.M. (2002). Artisan phakic iris claw intraocular lens for high primary and secondary hyperopia. *J Refract Surg*, 18, 697-707.
- Arne, J.L. & Lesueur, L.C. (2000). Phakic posterior chamber lenses for high myopia, functional and anatomical outcomes. *J Cataract Refract Surg*, 26, 369-374.
- Almustanyir, A. (2014). Computerized and Non-Computerized Colour Vision Tests. Master Thesis, University of Waterloo.
- American Optometric Association (AOA). (1990). Guidelines for the Use of Contact Lenses in Industrial Environments (position paper). Alexandria, VA.
- ANSI Z80.21. (2020). American National Standard for Ophthalmics: Instruments - General - Purpose Clinical Visual Acuity Charts. Alexandria, VA: The Vision Council.
- Autonomous Technologies Corporation. (2000). FDA Approval Application. Retrieved from: <<http://www.fda.gov/cdrh/pdf/p970043s005.pdf>>.
- Bailey, I.L., Bullimore, M.A., Raasch, T.W. & Taylor, H.R. (1991). Clinical grading and the effects of scaling. *Invest Ophthalmol Vis Sci*, 32, 422-432.
- Bailey, J.E. (1991). Color vision. In J.B. Eskridge, J.F. Amos & J.D. Bartlett (Eds.), *Clinical Procedures in Optometry* (pp.99-120). Philadelphia, PA: J.B. Lippincott.
- Barnett, M., Carrasquillo, K. G., & Schornack, M. M. (2020). Clinical Outcomes of Scleral Lens Fitting with a Data-driven, Quadrant-specific Design: Multicenter Review. *Optometry and Vision Science*, Pre-Publication.

Barnett, M., Ross, J., and Durbin-Johnson, B. Preliminary Clinical Exploration of Scleral Lens Performance on Normal Eyes. JCLRS [Internet]. 22 Nov 2018 [cited 2 Dec 2018]; 2(2):e14-e21.

Bausch & Lomb Surgical, Inc. (2000). FDA Approval Application. Retrived from: <<http://www.fda.gov/cdrh/pdf/p990027b.pdf>>.

Beauchamp, G.R., Black, B.C., Coats, D.K., Enzenauer, R.W., Hutchinson, A.K., Saunders, R.A., Simon, J.W., Stager, D.R., Stager, Jr DR., Wilson, M.E., Zabal-Ratner, J. & Felius, J. (2003). The management of strabismus in adults, clinical characteristics and treatment. *Journal of American Association for Pediatric Ophthalmology and Strabismus*, 7, 233-240.

Bedi, R., Touboul, D., Pinsard, L. & Colin, J. (2012). Refractive and topographic stability of Intacs in eyes with progressive keratoconus, five-year follow up. *J Refract Surg*, 28, 392-396

Borish, I.M. 1970. *Visual Acuity. Clinical Refraction*, 3rd ed. Boston, MA: Butterworth-Heinemann.

Bowden, T.J., Barnett, M. History of Scleral Lenses. In Barnett M, Johns LK. Eds, *Contemporary Scleral Lenses: Theory and Application*. (Chapter 1, 1-47) Bentham Science.

Brandl, S.G. (1996). In the line of duty: A descriptive analysis of police assaults and accident. *J Criminal Justice*, 24, 255-264.

Brandl, S.G. & Stroshine, M.S. (2012). The physical hazards of police work revisited. *Police Quarterly*, 15, 262-282.

Briggs, R., (1984) POST Study of Vision-Related Characteristics/Experiences, Unpublished report.  
Broome, P.W., & Classe, J.G. (1979). Long-term success in contact lens wear. *Contact Lens Forum*, (September), 15-27.

Bullimore, M.A., Bailey, I.L. & Wacker, R.T. (1991). Face recognition in age related maculopathy. *Invest Ophthalmol Vis Sci*, 32, 2020-2029.

Campbell, F.W., & Robson, J.G. (1968) Application of Fourier analysis to the visibility of gratings. *J Physiol*, 197:551-556.

Carmean, G., Chase, W. & Padgett, V.R. (2000). *Empirical Validation of Police Officer Visual Acuity Requirements*. Ontario, CA: MED-TOX Health Services Technical Report.

Cole, B. (1993). Does defective colour vision really matter? In B. Drum (Ed.), *Colour Vision Deficiencies XI* (Vol. 56, pp. 67-86): Springer Netherlands.

Cole, B.L. & Vingrys, A.J. (1983). Who fails lantern tests? *Doc Ophthalmol*, 55, 157-175.

Cole, B.L. & Maddocks, J.D. (1997). Defective colour vision is a risk factor in driving. In, Cavonius CR ed. *13th Symposium of the Internation Research Group on Colour Vision Deficiencies*. Pau France, Kluwer Acedemic, p 471-481.

Cole, B.L., Lian, K.Y. & Lakkis, C. (2006). The new Richmond HRR pseudoisochromatic test for colour vision is better than the Ishihara test. *Clin Exp Optom*, 89, 73-80.

Cole, B.L., Lian, K.Y., Sharpe, K. & Lakkis, C. (2006). Categorical color naming of surface color codes by people with abnormal color vision. *Optom Vis Sci*, 83, 879-886.

Cole, B.L., Lian, K.Y. & Lakkis, C. (2007). Can color vision defective subjects who pass the Farnsworth lantern test recognize surface color codes? *Aviat Space Environ Med*, 78, 21-25.



- Cole, B.L., Lian, K.Y. & Lakkis, C. (2008). Color vision assessment by Farnsworth lantern, results using alternative pass-fail criteria. *Aviat Space Environ Med* 2008, 79, 509-513.
- Cole, B.L. & Orenstein, J. (2003). Does the Farnsworth D-15 test predict the ability to name colors? *Clinical Exp Optom*, 86, 221-229.
- Cole, B.L., Lian, K.Y. & Lakkis, C. (2007). Using clinical tests of colour vision to predict the ability of colour vision deficient patients to name surface colours. *Ophthalmic Physiol Opt*, 27, 381-388.
- Cross, J.F., Cross, J. & Daly, J. (1971). Sex, race, age, and beauty as factors in recognition of faces. *Perception and Psychophysics*, 10, 393-396.
- Dain, S.J., Honson, V.J. & Curtis, C. (1993). Suitability of fluorescent tube light sources for the Ishihara test as determined by colorimetric methods. In, Drum B ed. *Color Vision Deficiencies XI*. Netherlands, Kluwer, 327-333.
- Dain, S. J., Atchison, D. A., Hovis, J. K., & Boon, M. Y. (2020). Lighting for color vision examination in the era of LEDs: the FM100Hue Test. *J Opt Soc Am A Opt Image Sci Vis*, 37(4), A122-A132.
- Davis, L.J., Schechtman, K.B., Wilson, B.S., Rosenstiel, C.E., Riley, C.H., Libassi, D.P., . . . Zadnik, K. (2006). Longitudinal changes in visual acuity in keratoconus. *Investigative Ophthalmology & Visual Science*, 47, 489-500
- Dennis, R.J., Green, R.P. & Ketchum, N.S. (1992). Clinical Medicine: The Medical Acceptability of Soft Contact Lens Wear by USAF Tactical Aircrews. *Aviat Space Environ Med*, 63, 122 127.
- Dennis, R.J., Apsey, D.A. & Ivan, D.J. (1993). Aircrew soft contact lens wear: A survey of USAF eyecare professionals. *Aviat Space Environ Med*, 64, 1044 1047.
- Dennis, R.J., Tredici, T.J., Ivan, D.J. & Jackson, W.G. (1996). The USAF aircrew medical contact lens study group: operational problems. *Aviat Space Environ Med*, 67, 303 307.
- Dirani, M., Couper, T., Yau, J., Ang, E.K., Islam, F.M.A., Snibson, G.R., Vajpayee, R.B. & Baird PN. (2010). Long term refractive outcomes and stability after excimer laser surgery for myopia. *J Cataract Refract Surg*, 36, 1709-1717.
- Dodson, S.R. 1993. Sighting-in on visual acuity. *Law and Order*. (June), 54-58.
- Dumbleton, K., Woods, C.A., Jones, L.W. & Fonn, D. (2013). The impact of contemporary contact lenses on contact lens discontinuation. *Eye & Contact Lens*, 39, 93-99
- Ellis, H.D. Shepherd, J., & Bruce A. (1973). The effects of age and sex upon adolescents' recognition of faces. *J Gen Psychol*, 123, 173.
- Espinosa-Mattar, Z., Gomez-Bastar, A., Graue-Hernandez, E. O., & Navas, A. (2012). DSAEK for implantable collamer lens dislocation and corneal decompensation 6 years after implantation. *Ophthalmic Surgery, Lasers and Imaging Retina*, 43 Online, e68-72
- European Committee for Standardisation (2009) ISO 8596:2009 Ophthalmic Optics. Visual Acuity Testing. Standard Optotype and its Presentation Beuth Verlag, Berlin.

- Farnsworth., D. 1947. The Farnsworth Dichotomous Test for Color Blindness Panel D-15 Manual. New York: The Psychological Corporation.
- Fernández-Velázquez, F. J. (2019). Performance and predictability of a new large diameter contact lens design in keratoconic corneae. *Contact Lens and Anterior Eye*, 42(3), 289-294
- Ferrari, G., & Rama, P. (2020). The keratoconus enigma: A review with emphasis on pathogenesis. *The Ocular Surface*, 18(3), 363-373.
- Ferris, F.L. Kassoff, A. Bresnick, G.H., & Bailey, I. (1982). New visual acuity charts for clinical research. *Am J Ophthalmol*, 94, 91-96.
- Fletcher, R.H., Fletcher, S.W. & Wagner, E.H. (1982). *Clinical Epidemiology- the Essentials*. Baltimore, MD: Williams and Wilkins.
- Fogt, J. S., Karres, M., & Barr, J. T. (2020). Changes in Symptoms of Midday Fogging with a Novel Scleral Contact Lens Filling Solution. *Optometry and Vision Science*, *Pre-Publication*.
- Freytag, E. & Sachs, J.C. (1969). Abnormalities of the central visual pathways contributing to seven Maryland traffic accidents. *JAMA*, 204, 119.
- Fuller, D. G., & Wang, Y. (2020). Safety and Efficacy of Scleral Lenses for Keratoconus. *Optom Vis Sci*, 97(9), 741-748.
- Ganesh, S., Brar, S., & Patel, U. (2018). Comparison of ReLEx SMILE and PRK in terms of visual and refractive outcomes for the correction of low myopia. *International Ophthalmology*, 38(3), 1147-1154.
- Giannoni, B. (1981). Entry-Level Vision Requirements Validation Study. Personnel Bureau, California Highway Patrol.
- Gibbs, G.H. (2001). The relationship between objective measures and subjective confidence in task performance with degraded visual acuity. In. *School of Kinesiology, Simon Fraser*, 112.
- Gimbel, H. V., LeClair, B. M., Jabo, B., & Marzouk, H. (2018). Incidence of implantable collamer lens-induced cataract. *Canadian Journal of Ophthalmology*, 53, 518-522.
- Ginsberg, A.P., Evans, D.W., Sekuler, R. & Harp, S.A. (1982). Contrast sensitivity predicts pilots' performance in aircraft simulators. *Am J Optom Physiol Opt*, 59, 105-109.
- Godefrooij, D.A., Gans, R. , Imhof, S.M. and Wisse, R.P. (2016), Nationwide reduction in the number of corneal transplantations for keratoconus following the implementation of cross-linking. *Acta Ophthalmol*, 94: 675-678.
- Goldberg, R.L. (1993). Eye Injuries Among LAPD Field Officers, 1987-1990. Los Angeles Police Department. Unpublished data.
- Goldberg, R.L. (1994). Field Testing of LAPD applicants who Pass the D-15. Los Angeles Police Department. Unpublished data.
- Goldberg, R.L. & Bible, M. (1993). Visual Testing of Incumbent Firefighters in the LAFD. Los Angeles Fire Department. Unpublished data.

- Goldberg, R.L. (2004). Development of a Short, Standardized and Objective Test of Color Vision Deficiency. County of Los Angeles. Unpublished data.
- Good, G.W. & Augsburger, A.R. (1987). Uncorrected visual acuity standards for police applicants. *J Police Sci Admin*, 15, 18-23.
- Good, G.W., Maisel, S.C. & Kriska, S.D. (1998). Setting an uncorrected visual acuity standard for police officer applicants. *J Appl Psychol*, 83, 817-824.
- Good, G.W., Fogt, N., Daum, K.M., Lynn & Mitchell, G. (2005). Dynamic visual fields of one eyed observers. *Optometry*, 76, 285-292.
- Gordon, M. O., Steger-May, K., Szczotka-Flynn, L., Riley, C., Joslin, C. E., Weissman, B. A., . . . Zadnik, K. (2006). Baseline factors predictive of incident penetrating keratoplasty in keratoconus. *American Journal of Ophthalmology*, 142(6), 923-930.
- Groome, W.R. & Johnson, C.A. (1993). Final Report on Entry Level Drawbridge Operator Vision Standard for the California Department of Transportation. Sacramento, CA: State Personnel Board.
- Groves, D. (1996). Refractive surgery: Defining rupture risks. *Phys Sports Med*, 24(5), 39-40.
- Hackman, R.J. & Holtzman, G.L. (1992). Color vision testing for the U.S. Naval Academy. *Mil Med*, 157(12), 651-657.
- Han, H.S. Song, J.S., & Kim, H.M. (2000). Long-term results of laser in situ keratomileusis for high myopia. *Korean J Ophthalmol*, 14(1), 1-6.
- Harbiyeli, I. I., Erdem, E., Isik, P., Yagmur, M., & Ersoz, R. (2020). Use of new-generation hybrid contact lenses for managing challenging corneas. *European Journal of Ophthalmology*, 112067212094275. doi:10.1177/1120672120942757
- Harthan, J., Nau, C. B., Barr, J., Nau, A., Shorter, E., Chimato, N. T., . . . Schornack, M. M. (2018). Scleral Lens Prescription and Management Practices: The SCOPE Study. *Eye Contact Lens*, 44, S228-S232.
- Hashemi, H., Asgari, S., Mehravaran, S., Emamian, M. H., & Fotouhi, A. (2020). Keratoconus after 40 years of age: a longitudinal comparative population-based study. *International ophthalmology*, 40(3), 583-589.
- Hatt, S.R., Leske, D.A., Kirgis, P.A., Bradley, E.A. & Holmes, J.M. (2007). The effects of strabismus on quality of life in adults. *Am J Ophthalmol*, 144, 643-647.
- Hazel, C.A. & Elliott, D.B. (2002). The dependency of logMAR visual acuity measurements on chart design and scoring rule. *Optom Vis Sci*, 79, 788-792.
- Hedin, A. & Lovsund, P. (1987). Effects of visual field defects on driving performance. *Documenta Ophthalmol Proc Series*, 49, 541-547.
- Holden, R.N. (1993). Eyesight standards - correcting myths. *FBI Law Enforcement Bulletin*, June, 1 6.
- Holt, D.G., Sikder, S. & Mifflin, M.D. (2012). Surgical management of traumatic LASIK flap dislocation with macrostriae and epithelial ingrowth 14 years postoperatively. *J Cataract Refract Surg*, 38, 357-361.

- Hovis, J.K. (1997). Long wavelength pass filters designed for the management of color vision deficiencies. *Optom Vis Sci*, 74, 222-230.
- Hovis, J.K. & Neumann, P. (1995). Colorimetric analyses of various light sources for the D-15 color vision test. *Optom Vis Sci*, 72, 667-678.
- Hovis, J.K. & Ramaswamy, S. (2006). Visual function of police officers who have undergone refractive surgery. *Am J Ind Med*, 49, 885-894.
- Hovis, J.K., Shaw, J. & Robinson, C.M. (1998). Spectacle and contact lens wear among police officers in Ontario. *Can J Optom*, 60, 24-32.
- Hovis, J.K. (2008). A review of the Constable Selection System Vision Screening Protocol. Prepared for Carroll Robinson, Community Safety and Correctional Services Selection Systems and Appointments Unit.
- Hovis, J.K., Lu, B. & Neumann, P. (1994). The ability of colour vision tests to predict performance in the identification of wire colour. *Can J Optom*, 55, 219-227.
- Implantable Contact Lens in Treatment of Myopia (ITM) Study Group (2003). U.S. Food and Drug Administration clinical trial of the Implantable Contact Lens for moderate to high myopia. *Ophthalmology*, 110, 255-266.
- Iskander, N.G., Peters, N.T., Anderson, P.E., & Gimbel, H.V (2001). Late traumatic flap dislocation after laser in situ keratomileusis. *J Cataract Refract Surg*, 27, 1111-1114.
- Jacobs, R.J., Hendicott, P.L., Murphy, B., Poppelwell, D. & Turner, P.J. (1998). Visual performance requirements for post PRK police recruits. *Clin Exper Opt*, 81, 163 172.
- Johnson, C.A. Casson, E.J., & Zadnik, K. (1992). Final report onet Entry-level Correctional Officer Vision Standards. Sacramento, CA: State of California Department of Corrections.
- Johnson, C.A. & Brintz, N. (1996). California Department of the Youth Authority Vision Study: Final Report. Entry-level vision standards for parole agents. Sacramento, CA: State of California Youth Authority.
- Johnson, C.A. & Keltner, J.L. (1983). Incidence of visual field loss in 20,000 eyes and its relationship to driving performance. *Arch Ophthalmol*, 101, 371-375.
- Johnson., L.N. & Baloh, F.G. (1991). The accuracy of confrontation visual field test in comparison with automated perimetry. (Vol.83) *J Ntl Med Assoc*, 10, 895-898.
- Kanakamedala, A., Salazar, H., Campagna, G., Chamberlain, P., Vadapalli, S., Orengo-Nania, S., & Khandelwal, S. (2019). Outcomes of Scleral Contact Lens Use in Veteran Population. *Eye & Contact Lens: Science & Clinical Practice*, Publish Ahead of Print.
- Kassar, B.S., Dresner, S.C., May, J.G., Marx, M.S. & Safir, A. (1984). Evaluation of the X Chrom lens and color deficiency. *CLAO Journal*, 10, 100-103.
- Kasneji, E., Sippel, K., Aehling, K., Heister, M., Rosenstiel, W., Schiefer, U. & Papageorgiou, E. (2014). Driving with binocular visual field loss, A study on a supervised on-road parcours with simultaneous eye and head tracking. *PLoS One*, 9, e87470.

- Keeney, A.H. (1968). Ophthalmic pathology in driver limitation. *Trans Am Acad Ophthalmol Otolaryngol*, 72, 737.
- Kennedy, R.H., Bourne, W.M., & Dyer, J.A. (1986). A 48-year clinical and epidemiologic study of keratoconus. *Am J Ophthalmol*, 101(3), 267-273
- Key, J.E. (1990). Are hard lenses superior to soft? *Cornea*, 9 (Suppl 1), S9-11.
- Kite, C.R. & King, J.N. (1961). A survey of the factors limiting the visual fields of motor vehicle drivers in relation to minimum visual fields and visibility standards. *Br J Physiol Opt*, 18, 85.
- Knorz, M.C., Jendritza, B., Liermann, A. Hugger, P., & Liesenhoff, H. (1998). LASIK for myopia correction, 2 year follow up. *Ophthalmol*, 95, 494-8.
- Koffler, B.H. & Sears, J.J. (2013). Myopia control in children through refractive therapy gas permeable contact lenses, Is it for real? *Am J Ophthalmol*, 156, 1076-1081
- Kok-van Aalphen, C.C. van der Linden, J.W., Visser, R., & Boh, A.H. (1985). Protection of the police against tear gas with soft lenses. *Mil Med*, 150, 451-454.
- Kong, J., Qin, X.J., Li, X.Y., Zhang, J.S. & Qu, B. (2010). Implantable collamer lens dislocation. *Ophthalmol*, 117, 399.
- Koppen, C., Kreps, E.O., Anthonissen, L., Van Hoey, M., Dhubhghaill, S.N., and Vermeulen, L. Scleral Lenses Reduce the Need for Corneal Transplants in Severe Keratoconus. *Am J Ophthalmol*. 2018 Jan;185:43-47. doi: 10.1016/j.ajo.2017.10.022. Epub 2017 Nov 16. PMID: 29103959.
- Kruk, R. & Regan, D. (1983). Visual test results compared with flying performance in telemetry tracked aircraft. *Aviation, Space, and Environmental Medicine*, 54, 906 911.
- Kushner, B.J. (1994). Binocular field expansion in adults after surgery for esotropia. *Arch Ophthalmol*, 112, 639-643.
- Kuyk, T.K., Veres, J.G. 3<sup>rd</sup>, Lahey, M.A., & Clark, D.J. (1986). The ability of protan color defectives to perform color dependant air traffic control tasks. *Am J Optom Physiol Opt*, 63, 582-586.
- Kuyk, T.K., Veres, J.G. 3<sup>rd</sup>, Lahey, M.A., & Clark, D.J. (1987). Ability of deutan color defectives to perform simulated air traffic control tasks. *Am J Optom Physiol Opt*, 64, 2-10.
- Lackner, B., Pieh, S., Schmidinger, G., Hanselmayer, G., Dejaco-Ruhswurm, I., Funovics, M.A. & Skorpik, C. (2003). Outcome after treatment of ametropia with implantable contact lenses. *Ophthalmol*, 110, 2153-2161.
- Lackner, B., Pieh, S., Schmidinger, G., Simader., C., Franz., C., Dejaco-Ruhswurm, I. & Skorpik, C. (2004). Long term results of implantation of phakic posterior chamber intraocular lenses. *J Cataract Refract Surg*, 30, 2269-2276.
- Levinger, S. & Pokroy, R. (2005). Keratoconus managed with intacs, One year results. *Arch of Ophthalmol*, 123, 1308-1314.

- Ling, J.J., Mian, S.I., Stein, J.D., Rahman, M., Poliskey, J., and Woodward, M.A. Impact of Scleral Contact Lens Use on the Rate of Corneal Transplantation for Keratoconus. *Cornea*. 2020 May 25. doi: 10.1097/ICO.0000000000002388. Epub ahead of print. PMID: 32452985.
- Maguen, E., Martinez, M., Rosner, I. R., Caroline, P., Macy, J., & Nesburn, A. B. (1991). The Use of Saturn II Lenses in Keratoconus. *Eye Contact Lens*, 17(1), 41-43
- Maguen, E., Caroline, P., Rosner, I. R., Macy, J. I., & Nesburn, A. B. (1992). The Use of the SoftPerm Lens for the Correction of Irregular Astigmatism. *Eye Contact Lens*, 18(3), 173-176
- Mahon, L.E. & Jacobs, R.J. (1991). Electronic flight information system displays and colour defective observers. *Clinical Exp Optom*, 74, 196-203.
- Maloney, R.K., Nguyen., L.H. & John, M.E. (2002). Artisan phakic intraocular lens for myopia, short term results of a prospective multicenter study. *Ophthalmol*, 109, 1631-1641.
- Mancuso, R. (1987). Survey of Myopic LAPD Officers. Los Angeles Police Department. Unpublished study.
- Mason, S.E. (1986). Age and gender as factors in facial recognition and identification. *Experimental Aging Research*, 12(3), 151-154.
- Matsumoto, E.R. Johnson, C.A., & Post, R.B. (1983). Effect of X chrom lens on chromatic discrimination and stereopsis in color deficient observers. *Am J Optom Physiol Opt*, 60, 279-302.
- McCulloch, D. L., Loffler, G., Colquhoun, K., Bruce, N., Dutton, G. N. & Bach, M. (2011). The effects of visual degradation on face discrimination. *Ophthalmic and Physiol Opt*, 31(3), 240-248.
- McDonald, M.B., Carr, J.D., Frantz, J.M., Kozarsky, A.M., Maguen, E., Nesburn, A.B., Rabinowitz, Y.S., Salz, J.J., Stulting, R.D., Thompson, K.P., & Waring, G.O. 3<sup>rd</sup>.(2001). Laser in situ keratomileusis for myopia up to -11 diopters with up to -5 diopters of astigmatism with the Summit Autonomous LADAR Vision Excimer Laser System. *Ophthalmol*, 108, 309-316.
- McDonnell, P.J., Nizam, A., Lynn, M.J., & Waring, G.O. 3<sup>rd</sup>. (1996). Morning to evening change in refraction, corneal curvature, and visual acuity 11 years after radial keratotomy in the PERK study. *Ophthalmol*, 103, 233-239.
- Menezo, J.L., Peris Martinez, C., Cisneros Lanuza, A.L. & Martinez-Costa, R. (2004). Rate of cataract formation in 343 highly myopic eyes after implantation of three types of phakic intraocular lenses. *J Refract Surg*, 20, 317-324.
- Miller, R.E., Thompson, W.T., Baldwin, J.B., Ivan, D.J., Tutt, R.C. & Hiers, P.L. (2005). Prospective Evaluation of Mesopic Night Vision and Night Vision Goggle Visual Acuity After Photorefractive Keratectomy (PRK). In. Brooks City Base TX, USAF School of Aerospace Medicine, p. 41.
- Moreland, J.D, Westland, S., Cheung, V. & Dain, S.J. (2010). Quantitative assessment of commercial filter "aids" for red green colour defectives. *Ophthalmic Physiol Opt*, 30, 685-692.
- Moshirfar, M., Dodd, J.G., Muir, G.J. & Gagnon, M.R. (2007a). Occurrence and prognosis of non penetrating injuries in eyes with Verisyse iris supported phakic intraocular lens. *J Cataract Refract Surg*, 33, 166-169.

Moshirfar, M., Holz, H.A. & Davis, D.K. (2007b). Two year follow up of the Artisan/Verisyse iris supported phakic intraocular lens for the correction of high myopia. *J Cataract Refract Surg*, 33, 1392-1397.

Moshirfar, M., Stagg, B. C., Muthappan, V., & Vasavada, S. A. (2014). Traumatic dislocation of implanted collamer phakic lens: a case report and review of the literature. *The Open Ophthalmology Journal*, 8, 24-26.

Milburn, H.J. & Mertens, H.W. (1993). Validation of an inexpensive test illuminant for aeromedical color vision screening. DOT/FAA/AM 93/16. Oklahoma City, FAA Civil Aeromedical Inst. NTIS.

Munoz, G., Montes Mico, R., Belda, J.I. & Alió, J.L. (2003). Cataract after minor trauma in a young patient with an iris fixated intraocular lens for high myopia. *Am J Ophthalmol*, 135, 890-891.

Nau, A. C. (2008). A Comparison of Synergeyes Versus Traditional Rigid Gas Permeable Lens Designs for Patients With Irregular Corneas. *Eye & Contact Lens: Science & Clinical Practice*, 34(4), 198-200.

Nichols, J. J., & Starcher, L. (2020). Contact Lenses 2019. *Contact Lens Spectrum*, 35, 18-25.

Ng, J. S., Self, E., Vanston, J. E., Nguyen, A. L., & Crognale, M. A. (2015). Evaluation of the Waggoner Computerized Color Vision Test. *Optometry & Vision Science*, 92, 480-486.

Ng, J. S., & Morton, W. A. (2018a). Case Report: Invalidation of the Farnsworth D15 Test in Dichromacy Secondary to Practice. *Optom Vis Sci*, 95, 272-274.

Ng, J. S., & Liem, S. C. (2018b). Can the Farnsworth D15 Color Vision Test Be Defeated through Practice? *Optom Vis Sci*, 95, 452-456.

Nidek Technologies Inc. (2000). FDA Approval Application. Retrieved from: <<http://www.fda.gov/cdrh/pdf/p970053s002b.pdf>>.

Nilsson, S.E.G. Lövsund, P., & Oberg, P.A.. (1981). Contact lenses and mechanical trauma to the eye. *Acta Ophthalmol*, 59, 402-408.

Nilsson, S.E.G. & Andersson, L. (1982). The use of contact lenses in environments with organic solvents, acids or alkalis. *Acta Ophthalmol*, 60, 599-608.

Nilsson, S. & Lindh, H. (1984). Daily contact lens wear, a three year follow-up. *Acta Ophthalmol*, 62, 556-565.

Nilsson, S. & Persson, G. (1986). Low complication rate in extended wear contact lenses. *Acta Ophthalmol*, 64, 88-92.

Norman, L.G. (1980). Medical aspects of road safety. *Lancet*, 1, 1039-1045.

O'Donnell, C., & Efron, N. (2012). Diabetes and contact lens wear. *Clinical and Experimental Optometry*, 95, 328-337.

O'Neal, M.R. & Miller, R.E. (1988). Further investigations of contrast sensitivity and visual acuity in pilot detection of aircraft. In: Wright Patterson AFB Dayton, OH, USAF School of Aerospace Medicine Brooks AFB, TX.

Pate, A. & Hamilton, E. (1991). *The Big Six, Policing America's Largest Cities*. Washington, D.C: The Police Foundation.

Patel, C.K., Hanson, R., McDonald, B., & Cox, N. (2001). Case reports and small case series: Late dislocation of a lasik LASIK flap caused by a fingernail. *Arch Ophthalmol*, 119, 447-449.

Pecego, M., Barnett, M., Mannis, M. J., & Durbin-Johnson, B. (2012). Jupiter Scleral Lenses. *Eye & Contact Lens: Science & Clinical Practice*, 38(3), 179-182

Piñero, D.P., Carracedo, G., Ruiz-Fortes, P., and Pérez-Cambrodí, R.J. Comparative analysis of the visual performance and aberrometric outcomes with a new hybrid and two silicone hydrogel multifocal contact lenses: a pilot study. *Clin Exp Optom*. 2015 Sep;98(5):451-8. doi: 10.1111/cxo.12299. PMID: 26390908.

Pinizzotto, A.M. et al. (2012). Restraint in the Use of Deadly Force: A Preliminary Study. *FBI Law Enforcement Bulletin*, June, 2012, pp. 1-11.

Polse, K. et al. (1990). *Contact Lens Use Under Adverse Conditions - Applications in Military Aviation*. Washington DC: National Academy Press.

Pop, M. & Payette, Y. (2000). Photorefractive keratectomy versus laser in situ keratomileusis, a control matched study. *Ophthalmol*, 107, 251-257.

POST. 1984. *Color Vision for Patrol Officers*. Sacramento, CA: Commission on Peace Officer Standards and Training. Unpublished study.

Pritchard, N., Fonn, D. & Brazeau, D. (1999). Discontinuation of contact lens wear: a survey. *Int Contact Lens Clin*, 26, 157-162.

Pullum, K. W., Whiting, M. A., & Buckley, R. J. (2005). Scleral Contact Lenses: The Expanding Role. *Cornea*, 24(3), 269-277. doi:10.1097/01.icc.0000148311.94180

Quah, S.A. & Kaye, S.B. (2004). Binocular visual field changes after surgery in esotropic amblyopia. *Invest Ophthalmol & Vis Sci*, 45, 1817-1822.

Qasem, Q., Kirwan, C. & O'Keefe, M. (2010). 5 year prospective follow up of Artisan phakic intraocular lenses for the correction of myopia, hyperopia and astigmatism. *Ophthalmologica*, 224, 283-290.

Raasch, T.W., Bailey, I.L. & Bullimore, M.A. (1998). Repeatability of visual acuity measurement. *Optom Vis Sci*, 75, 342-348.

Rabin, J. (1994). Optical defocus: Differential effects on size and contrast recognition threshold. *Invest Ophthalmol Vis Sci*, 3, 646-648.

Ramaswamy, S. & Hovis, J.K. (2004). Ability of the D 15 panel tests and HRR pseudoisochromatic plates to predict performance in naming VDT colors. *Vis Neurosci*, 21, 455-460.

Rengstorff, R.H. & Black, C.J. (1974). Eye protection from contact lenses. *J Am Optom Assoc*, 45(3), 270-276.

Richdale, K., Sinnott, L.T., Skadahl, E. & Nichols, J. (2007). Frequency of and factors associated with contact lens dissatisfaction and discontinuation. *Cornea*, 26, 168-174.



- Robbins, J.C. (1977). A three-year retrospective soft lens contact lens study. In Proc 2nd Natl Res Symp Soft Contact Lenses Int. Congr. Ser. No. 398, 57-61. Excerpta Medica, Amsterdam.
- Royall, W.W. (1977). Soft contacts and law enforcement. *Contact Lens Forum*, 3, 15-17.
- Sanders, D.R. & Vukich, J.A., (2002). Incidence of lens opacities and clinically significant cataracts with the implantable contact lens: comparison of two lens designs. *J Refract Surg*, 18, 673-682.
- Sanders, D.R., Vukich, J.A., Doney, K. & Gaston, M. (2003). U.S. Food and Drug Administration clinical trial of the Implantable Contact Lens for moderate to high myopia. *Ophthalmol*, 110, 255- 266.
- Sanders, D.R., Doney, K. & Poco, M. (2004). United States Food and Drug Administration clinical trial of the Implantable Collamer Lens (ICL) for moderate to high myopia, three year follow up. *Ophthalmol*, 111, 1683-1692.
- Saraç, Ö., Kars, M. E., Temel, B., & Çağıl, N. (2019). Clinical evaluation of different types of contact lenses in keratoconus management. *Contact Lens and Anterior Eye*, 42(5), 482-486
- Sarikkola, A.-U., Sen, H. N., Uusitalo, R. J., & Laatikainen, L. (2005). Traumatic cataract and other adverse events with the implantable contact lens. *Journal of Cataract & Refractive Surgery*, 31), 511-524.
- Schade, O.H. (1956). Optical and photoelectronic analog of the eye. *J Opt Soc Am*, 46, 721 39
- Schanzlin, D.J., Santos, V.R., Waring, G.O. 3<sup>rd</sup>, Lynn, M., Bourque, L., Cantillo, N., Edwards, M.A., Justin, N., Reinig, J., & Roszka-Duggan V. (1986). Diurnal change in refraction, corneal curvature, visual acuity, and intraocular pressure after radial keratotomy in the PERK study. *Ophthalmol*, 93(2), 167-175.
- Schanzlin, D.J. (1999). Studies of intrastromal corneal ring segments for the correction of low to moderate myopic refractive errors. *Trans Am Ophthal Soc*, 97, 815-890.
- Schmitz, J.W., McEwan, G.C. & Hofmeister, E.M. (2012). Delayed presentation of traumatic dislocation of a Visian Implantable Collamer Lens. *J Refract Surg*, 28, 365-367.
- Schornack, M. (2017). Medical Indications for Scleral Lenses. In Barnett M, Johns LK. Eds, *Contemporary Scleral Lenses: Theory and Application*. (Chapter 5, 138-165) Bentham Science.
- Schornack, M., Nau, C., Nau, A., Harthan, J., Fogt, J., & Shorter, E. (2019). Visual and physiological outcomes of scleral lens wear. *Contact Lens and Anterior Eye*, 42(1), 3-8
- Shaw, J. & Gledhill, N. (1995). Constable Selection Project. Report on Job Review Phase. Medical, Physical, Skills and Abilities. Recommendations, Tests and Standards. Submitted to the Ministry of the Solicitor General and Correctional Services of Ontario.
- Sheedy, J.E. (1980). Police vision standards. *J Police Sci Admin*, 8(3), 275-285.
- Shojaei, A., Mohammad Rabei, H., Eslani, M., Elahi, B. & Noorizadeh F. (2009). Long term evaluation of complications and results of photorefractive keratectomy in myopia, an 8 year follow up. *Cornea*, 28, 304-310.
- Shinar, D. & Gilead, E. (1987). Contrast Sensitivity as a Predictor of Complex Target Detection. *Human Factors*, 31, 1194 1197.

- Siu, A.W. & Yap, M. (2003). The performance of color deficient individuals on airfield color tasks. *Aviat Space & Environ Med*, 74, 546-550.
- Spilberg, S.W. (1993). Analysis of LAPD Officer initiated Shootings by Distance and Time of Day. Sacramento, CA: State of California Commission on Police Officers Standards and Training. Unpublished data.
- Sridhar, M.S., Rapuano, C.J. & Cohen, E.J. (2001). Accidental self-removal of a flap--a rare complication of laser in situ keratomileusis surgery. *Am J Ophthalmol*, 132, 780-782.
- Stager, P. & Hameluck, D. (1986). Contrast sensitivity and visual detection in search and rescue. In. *Downsview, ON: Defence and Civil Institute of Environmental Medicine*, p. 41.
- Steward, J.M & Cole, B.L. (1989). What do color vision defectives say about everyday tasks? *Optom Vis Sci*, 66(5), 288-295.
- Suiter, B.G., Twa, M.D., Ruckhofer, J. & Schanzlin, D.J. (2000). A comparison of visual acuity, predictability, and visual function outcomes after intracorneal ring segments and laser in situ keratomileusis. *Trans Am Ophthalmol Soc*, 98, 51-55.
- Sulley, A., Young, G., & Hunt, C. (2017). Factors in the success of new contact lens wearers. *Contact Lens and Anterior Eye*, 40(1), 15-24.
- Sulley, A., Young, G., Hunt, C., McCreedy, S., Targett, M.-T., & Craven, R. (2018). Retention Rates in New Contact Lens Wearers. *Eye & Contact Lens: Science & Clinical Practice*, 44, S273-S282.
- Summit Technologies Inc. (1999). FDA Approval Application, Retrieved from: <[www.fda.gov/cdrh/pdf/p930034s013b.pdf](http://www.fda.gov/cdrh/pdf/p930034s013b.pdf)>.
- Swarbrick, H.A., Nguyen, P., Nguyen, T. & Pham, P. (2001). The ChromaGen contact lens system, colour vision test results and subjective responses. *Ophthalmic Physiol Opt*, 21, 182-196.
- Swarbrick, H.A. (2006). Orthokeratology review and update. *Clinical and Experimental Optometry*, 89, 124-143.
- Temme, L.A., Ricks, E., Morris, A. & Sherry, D. (1991). Visual contrast sensitivity of U.S. Navy jet pilots. *Aviat Space Environ Med*, 62, 1032-1036.
- Uçakhan, Ö. Ö., & Yeşiltaş, Y. S. (2020). Correction of Irregular Astigmatism With New-Generation Hybrid Contact Lenses. *Eye & Contact Lens: Science & Clinical Practice*, 46(2), 91-98.
- Verriest, G., Neubauer, D., Marre, M., & Uvijls, A. (1980). New investigations concerning the relationships between congenital color vision defects and road traffic security. In G. Verriest, ed. *Colour Vision Deficiencies*. V. Bristol: Hilger, 331-342.
- Vinger, P.F. et al. (1996). Ruptured globes following radial and hexagonal keratotomy surgery. *Arch Ophthalmol*, 114, 129-134.
- Vingrys, A.J. (2002). The case against protan drivers holding professional driving licenses. *Clinical and Experimental Optometry*, 85, 46-48.

- Von Noorden, G.K. (1990). *Binocular Vision & Ocular Motility, Theory & Management of Strabismus*. St. Louis, MO: Mosby Yr Bk.
- Wagner, H., Barr, J.T., & Zadnik, K. (2007). Collaborative Longitudinal Evaluation of Keratoconus (CLEK) Study: Methods and findings to date. *Contact Lens and Anterior Eye*, 30(4), 223-232.
- Walker, M. K., Bergmanson, J. P., Miller, W. L., Marsack, J. D., & Johnson, L. A. (2016). Complications and fitting challenges associated with scleral contact lenses: A review. *Contact Lens and Anterior Eye*, 39(2), 88-96.
- Waring, G.O. & Durrie, D.S. (2008). Emerging trends for procedure selection in contemporary refractive surgery, consecutive review of 200 cases from a single center. *J Refract Surg*, 24(4), S419-23.
- Wells, G.A., Brown, J.J., Casson, E.J., Easterbrook, M. & Trottier, A.J. (1997). To wear or not to wear, current contact lens use in the Royal Canadian Mounted Police. *Can J Ophthalmol*, 32, 158-162.
- Wilson, H. R., Loffler, G. & Wilkinson., F. (2002). Synthetic faces, face cubes, and the geometry of face space. *Vis Research*, 42, 2909-2923.
- Wolfe, R.J.B., Vingrys, A.J. & Cole, B.L. (2002). Protans and driving safety. *Clinical and Experimental Optometry*, 85, 399-402.
- Wood, J.M., McGwin, G., Elgin, J., Vaphiades, M.S., Braswell, R.A., DeCarlo, D.K., Kline, L.B., Meek, G.C., Searcey, K., & Owsley, C. (2009). On road driving performance by persons with hemianopia and quadrantanopia. *Invest Ophthalmol Vis Sci*, 50, 577-585.
- Yoon, H., Macaluso, D.C., Moshirfar, M. & Lundergan, M. (2002). Traumatic dislocation of an Ophtec Artisan phakic intraocular lens. *J Refract Surg*, 18, 481-483.
- Young, G., Veys, J., Pritchard, N., Coleman, S. (2002). A multi centre study of lapsed contact lens wearers. *Ophthalmic Physiol Opt*, 22, 516-527.
- Zisman, F. & Adams, A.J. (1985). *Color Vision Screening Policy*. (September 11). Berkeley, CA: University of California, School of Optometry.

